arXiv:1504.03448v3 [math.AG] 21 Jun 2016

TORIC NETWORKS, GEOMETRIC R-MATRICES AND GENERALIZED DISCRETE TODA LATTICES

REI INOUE, THOMAS LAM, AND PAVLO PYLYAVSKYY

Abstract. We use the combinatorics of toric networks and the double affine geometric R-matrix to define a three-parameter family of generalizations of the discrete Toda lattice. We construct the integrals of motion and a spectral map for this system. The family of commuting time evolutions arising from the action of the R-matrix is explicitly linearized on the Jacobian of the spectral curve. The solution to the initial value problem is constructed using Riemann theta functions.

1. Introduction

We study a generalization of the discrete Toda lattice parametrized by a triple of integers (n, m, k), which corresponds to a network on a torus with n horizontal wires, m vertical wires and k *shifts* at the horizontal boundary. An example of the kind of toric network that we consider is given in Figure 1.



Figure 1. The toric network for $n = 3$, $m = 4$ and $k = 0$.

The phase space $\mathcal{M}$ of our system is the space of parameters $q_{ij} \in \mathbb{C}$ assigned to each of the crossings of the wires (a $3 \times 4 = 12$ dimensional space for the example in Figure 1). We construct two families of commuting discrete time evolutions acting on the parameters q_{ij}, together generating an action of $\mathbb{Z}^m \times \mathbb{Z}^N$, where $N := \gcd(n, k)$. These time evolutions act as birational transformations of the phase space. The purpose of this paper is to study the algebro-geometrical structure of these maps and to solve the corresponding initial value problem.

The rational transformations of the phase space come from the *affine geometric R-matrix*, acting on the parameters of two adjacent parallel wires, either horizontal or vertical. The affine geometric R-matrix arises in the theory of affine geometric crystals

---

Date: August 31, 2015 (Final version: June 19, 2016).

R.I. was partially supported by JSPS KAKENHI Grant Number 26400037.

T.L. was partially supported by NSF grants DMS-1160726, DMS-1464693, and a Simons Fellowship.

P. P. was partially supported by NSF grants DMS-1148634, DMS-1351590, and Sloan Fellowship.

[BK, KN1, KN2], being a birational lift of the *combinatorial R-matrix* of certain tensor products of Kirillov-Reshetikhin crystals for $U_q(\widehat{\mathfrak{sl}}_n)$ [KKMMNN]. Geometric R-matrices also arises independently in the study of Painlevé equations [KNY] and total positivity [LP13]; see also [Kir, Eti].

The geometric R-matrix satisfies the Yang-Baxter relation and we show in Theorem 2.4 that the action of the geometric R-matrix generates commuting birational actions of two affine Weyl groups W and $\tilde{W}$: one swapping horizontal wires, and one swapping vertical wires. The commutativity was proved in [KNY] for the case $k = 0$; we extend it here to arbitrary k.

The $\mathbb{Z}^m \times \mathbb{Z}^N$ time-evolutions of our dynamical system come from the subgroup of translation elements in the corresponding affine Weyl group. In the case $(n, m, k) = (n, 2, n-1)$, a rational map given by one of the $\mathbb{Z}^{m=2}$ actions corresponds to the discretization of the well-known n-periodic Toda lattice equation, studied in [HTI].

The dynamics that we study come from positive birational maps, and they tropicalize to the box-ball system [TS] and the combinatorics of jeu-de-taquin (see §2.5.2).

To study the dynamics of our generalized discrete Toda lattice, we construct a spectral map

$$\begin{aligned} \phi : \mathcal{M} &\longrightarrow \{\text{plane algebraic curve}\} \times \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A \\ (q_{ij}) &\longmapsto (C_f, \mathcal{D}, (c_1, \ldots, c_M), O, A) \end{aligned}$$

where C_f is a spectral curve, $\mathcal{D} \in \mathrm{Pic}^g(C_f)$ is degree g divisor, and the remaining data is explained in §5. We show that when appropriately restricted, the spectral map is an injection. Such spectral data is frequently encountered in the theory of integrable systems, and our approach follows that of van Moerbeke and Mumford [vMM]. Van Moerbeke and Mumford used similar spectral data to study periodic difference operators.

The heart of our work is the calculation of the double affine geometric R-matrix action in terms of the spectral data: we show that the translation subgroup $\mathbb{Z}^m \times \mathbb{Z}^N$ acts as constant motions on the Jacobian of C_f, while the symmetric subgroups of W and of $\tilde{W}$ act by permuting the additional data R_O and R_A (which are certain special points on C_f).

We explicitly invert (for $N = 1$) the spectral map ϕ using Riemann theta functions, and give a solution to the initial value problem. This extends work of Iwao [I08, I10], who studied the initial value problem of the $\mathbb{Z}^m$ action in the $(n, m, n-1)$ and $(n, m, 0)$ cases. We relate our theta function solutions to the octahedron recurrence [Spe] via Fay's trisecant identity [Fay]. We also give an interpretation of our dynamics in terms of dimer model transformations [GK].

Outline. The outline of this paper is as follows: in §2, we introduce a family of commuting actions on the toric network, generalizating the discrete Toda lattice. We briefly summarize the main results of this paper in §2.4, and give some examples and applications in §2.5. In §3, we introduce a space $\mathcal{L}$ of Lax matrices for our system, and construct the spectral curves C_f, where $f \in \mathbb{C}[x, y]$. We also study certain special points on our spectral curves, and analyze some of the singular points. In §5, we study the spectral map which sends a point in the phase space to the spectral curve C_f, a divisor $\mathcal{D}$ on

C_f, and some additional data. We modify the strategy in [vMM] to fit our situation. In §4 and §6 we present the proofs of results in §3 and §5 respectively. In particular, the coefficients of the spectral curves (which are integrals of motion) are described explicitly in terms of the combinatorics of these networks. In §7, we study the vertical $\mathbb{Z}^m$ actions, the horizontal $\mathbb{Z}^N$ actions, and the snake path actions. We show that all the actions preserve the spectral curve, and the commuting $m+N$ actions induce constant motions on the Picard group of C_f, through the map ϕ. In §8, we solve, for the case of $N=1$, the initial value problem for the commuting actions by constructing the inverse of ϕ in terms of the Riemann theta function. Our method extends the strategy in [I10]. For $N>1$, a solution is given that relies on a technical condition. In §9, we show that the theta function solution for the system satisfies the octahedron recurrence, by specializing Fay's trisecant identity for the Riemann theta function. In §10, we study the symmetry between the network and its transposition obtained by swapping the roles of the vertical and the horizontal wires. In §11, we realize the R-matrix transformation on a toric network in terms of transformations of the honeycomb dimer model on a torus [GK]. An explicit interpretation of the R-matrix as a cluster transformation will be the subject of future work [ILP], so we do not elaborate on the relation to cluster structures here.

Future directions. There are several systematic ways to construct integrable rational maps using combinatorial objects on surfaces, such as directed networks, electrical networks, and bipartite graphs [LP13, GK, GSTV]. In particular, the R-matrix of the present work has an "electrical" analogue [LP12b]. It would be interesting to study these discrete-time dynamical systems from the view point of algebraic geometry (Cf. [FM]). In all these cases, the rational transformations are additionally positive, and can be tropicalized to piecewise-linear maps that generate a discrete-time and discrete-space dynamical system. It is natural to consider the tropical counterpart of our work using tropical geometry as in [IT], where tropical curves and tropical theta functions are used to study the piecewise-linear map arising from the $(n,m,k)=(n,2,n-1)$ case.

Acknowledgments. We thank Rick Kenyon for helpful discussions. We also thank the anonymous referee for kind comments which improved the manuscript.

2. Dynamical system from a toric network

We study the action of the geometric R-matrix on an array of variables. The geometric R-matrix generates an action of a product of commuting (extended) affine symmetric groups, acting as birational transformations.

2.1. **The affine geometric R-matrix.** For a vector $\mathbf{a}=(a_1,\dots,a_n)$, let $\mathbf{a}^{(i)} := (a_{i+1},a_{i+2},\dots,a_n,a_1,\dots,a_i)$. Let $\mathbf{a}=(a_1,\dots,a_n)$ and $\mathbf{b}=(b_1,\dots,b_n)$ be two vectors. Define the *energy* $E(\mathbf{a},\mathbf{b})$ to be

$$E(\mathbf{a},\mathbf{b})=\sum_{i=0}^{n-1}\left(\prod_{j=1}^{i}b_j\prod_{j=i+2}^{n}a_j\right).$$

Define the affine geometric R-matrix to be the transformation $R : (\mathbf{a}, \mathbf{b}) \mapsto (\mathbf{b}', \mathbf{a}')$, given by

$$b_i' = b_i \frac{E(\mathbf{a}^{(i)}, \mathbf{b}^{(i)})}{E(\mathbf{a}^{(i-1)}, \mathbf{b}^{(i-1)})} \qquad \text{and} \qquad a_i' = a_i \frac{E(\mathbf{a}^{(i-1)}, \mathbf{b}^{(i-1)})}{E(\mathbf{a}^{(i)}, \mathbf{b}^{(i)})}.$$

We will usually think of R as a birational transformation of $\mathbb{C}^n \times \mathbb{C}^n$.

It is easy to see

$$\prod_{i=1}^{n} a_i = \prod_{i=1}^{n} a_i' \qquad \text{and} \qquad \prod_{i=1}^{n} b_i = \prod_{i=1}^{n} b_i'. \tag{2.1}$$

2.2. **Description of dynamics.** The R-matrix can be used to act on a rectangular array of variables, by acting on consecutive pairs of rows or of columns.

Let $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}$ be three positive integers, and let $0 \leq \mathfrak{d} < \mathfrak{c}$ be a nonnegative integer satisfying $\gcd(\mathfrak{c}, \mathfrak{d}) = 1$. Note that we allow $\mathfrak{d} = 0$ if and only if $\mathfrak{c} = 1$. Let $\mathfrak{d}^{-1}$ be the unique number in the range $0 \leq \mathfrak{d}^{-1} < \mathfrak{c}$ such that $\mathfrak{d}\mathfrak{d}^{-1} = 1 \mod \mathfrak{c}$. For a positive integer a, we define $[a] := \{1, 2, \ldots, a\}$.

We consider an array of $\mathfrak{a} \times \mathfrak{b} \times \mathfrak{c}$ variables $\{q_{a,b,c}\}_{a \in [\mathfrak{a}], b \in [\mathfrak{b}], c \in [\mathfrak{c}]}$. We consider two distinct ways to arrange them in a two-dimensional rectangular array:

$$q_{i,j} = q_{a,b,c} \text{ for } i = a,\ j = b + \mathfrak{b}(c-1) \tag{2.2}$$

and

$$\tilde{q}_{i,j} = q_{a,b,c} \text{ for } i = \mathfrak{b} + 1 - b,\ j = \mathfrak{a}\mathfrak{d}^{-1}c - a + 1 \mod \mathfrak{a}\mathfrak{c}. \tag{2.3}$$

By convention, $q_{i,j}$ denotes the entry in the *i-th column* and *j-th row*. We call these arrays Q and $\tilde{Q}$. The first array Q has dimensions $\mathfrak{b}\mathfrak{c} \times \mathfrak{a}$ and the second array $\tilde{Q}$ has dimensions $\mathfrak{a}\mathfrak{c} \times \mathfrak{b}$. Note that changing $\mathfrak{a} \mapsto \mathfrak{b}$, $\mathfrak{d} \mapsto \mathfrak{d}^{-1}$, $q_{a,b,c} \mapsto q_{1-b,1-a,\mathfrak{d}^{-1}c}$ swaps Q and $\tilde{Q}$.

Example 2.1. Let $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (2, 2, 2, 1)$. The two arrays are as follows.

$$Q := \begin{pmatrix} q_{1,1,1} & q_{2,1,1} \\ q_{1,2,1} & q_{2,2,1} \\ q_{1,1,2} & q_{2,1,2} \\ q_{1,2,2} & q_{2,2,2} \end{pmatrix}, \qquad \tilde{Q} := \begin{pmatrix} q_{2,2,1} & q_{2,1,1} \\ q_{1,2,1} & q_{1,1,1} \\ q_{2,2,2} & q_{2,1,2} \\ q_{1,2,2} & q_{1,1,2} \end{pmatrix}. \tag{2.4}$$

Example 2.2. Let $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (3, 2, 3, 2)$. The two arrays are as follows.

$$Q := \begin{pmatrix} q_{1,1,1} & q_{2,1,1} & q_{3,1,1} \\ q_{1,2,1} & q_{2,2,1} & q_{3,2,1} \\ q_{1,1,2} & q_{2,1,2} & q_{3,1,2} \\ q_{1,2,2} & q_{2,2,2} & q_{3,2,2} \\ q_{1,1,3} & q_{2,1,3} & q_{3,1,3} \\ q_{1,2,3} & q_{2,2,3} & q_{3,2,3} \end{pmatrix}, \qquad \tilde{Q} := \begin{pmatrix} q_{3,2,1} & q_{3,1,1} \\ q_{2,2,1} & q_{2,1,1} \\ q_{1,2,1} & q_{1,1,1} \\ q_{3,2,3} & q_{3,1,3} \\ q_{2,2,3} & q_{2,1,3} \\ q_{1,2,3} & q_{1,1,3} \\ q_{3,2,2} & q_{3,1,2} \\ q_{2,2,2} & q_{2,1,2} \\ q_{1,2,2} & q_{1,1,2} \end{pmatrix}. \tag{2.5}$$

The arrays Q and $\tilde{Q}$ correspond to a network on a torus in the following way. Let G a network embedded into a torus, as illustrated in Figure 2. Each rectangle of red lines denotes the fundamental domain of the torus. There are $\mathfrak{a}$ vertical "wires" (directed upwards), forming $\mathfrak{a}$ simple curves in the torus with the same homology class, and $\mathfrak{b}\mathfrak{c}$ horizontal wires (directed to the right), which form $\mathfrak{b}$ simple curves on the torus with another homology class: the top $\mathfrak{b}\mathfrak{d}$ right ends of horizontal wires cross the top edge and come out on the other side. We set q_{ij} at the crossing of the i-th vertical and the j-th horizontal wires, then Q corresponds to the configuration of the q_{ij} on G. When we see G from the inside of the torus, the roles of the vertical and horizontal wires are interchanged; there are $\mathfrak{b}$ vertical wires forming $\mathfrak{b}$ simple curves, and $\mathfrak{a}\mathfrak{c}$ horizontal wires forming $\mathfrak{a}$ simple curves. The top $\mathfrak{a}\mathfrak{d}^{-1}$ ends of horizontal wires cross the top edge and come out on the other side. The array $\tilde{Q}$ corresponds to this "from-inside" configuration, where $\tilde{q}_{ij}$ is placed at the crossing of the i-th vertical and the j-th horizontal wires.

See Figure 3 for the network in the case of Example 2.2, where the lines with the same color in the two pictures are identical simple curves in G.


1
2
⋯
$\mathfrak{a}$
1
2
⋮
$\mathfrak{b}\mathfrak{d}$
$\mathfrak{b}\mathfrak{d}+1$
⋮
$\mathfrak{b}\mathfrak{c}$
1
2
⋯
$\mathfrak{a}$
1
2
⋮
$\mathfrak{b}\mathfrak{d}$
$1+\mathfrak{b}(\mathfrak{c}-\mathfrak{d})$
$2+\mathfrak{b}(\mathfrak{c}-\mathfrak{d})$
⋮
$\mathfrak{b}\mathfrak{c}$
1
2
⋯
$\mathfrak{a}$
1
⋮
$\mathfrak{b}(\mathfrak{c}-\mathfrak{d})$

FIGURE 2. Toric network

Definition 2.3. Let $A = \{a_{i,j}\}_{i\in[s],j\in[r]}$ be an $r \times s$ array. Let $\mathbf{a}_i = (a_{i,j})_{j\in[r]}$, and write $\mathbf{a}_i^T$ for the transpose of $\mathbf{a}_i$. For $1 \le \ell \le s-1$, define the array $s_\ell(A)$ by

$$s_\ell(A) = \left(\mathbf{a}_1^T, \dots, \mathbf{a}_{\ell-1}^T, (\mathbf{a}'_{\ell+1})^T, (\mathbf{a}'_\ell)^T, \mathbf{a}_{\ell+2}^T, \dots, \mathbf{a}_s^T\right) \tag{2.6}$$

if $A = (\mathbf{a}_1^T, \dots, \mathbf{a}_{\ell-1}^T, \mathbf{a}_\ell^T, \mathbf{a}_{\ell+1}^T, \mathbf{a}_{\ell+2}^T, \dots, \mathbf{a}_s^T)$, and $R(\mathbf{a}_\ell, \mathbf{a}_{\ell+1}) = (\mathbf{a}'_{\ell+1}, \mathfrak{a}'_\ell)$. Also define an array $\pi(A)$ by the formula

$$\pi(A)_{i,j} = a_{i,j-1}. \tag{2.7}$$

Here we consider the first index modulo r and the second index modulo s.

Let $\mathcal{M} \simeq \mathbb{C}^{\mathfrak{a}\mathfrak{b}\mathfrak{c}}$ be the phase space of our dynamical system and regard $(q_{a,b,c})_{a\in[\mathfrak{a}],b\in[\mathfrak{b}],c\in[\mathfrak{c}]}$ as coordinates on $\mathcal{M}$. Applying the above definition to the array Q, we obtain operators $s_1, \dots, s_{\mathfrak{a}-1}$ and $\tilde{\pi}$ acting on $\mathcal{M}$. Similarly, applying this definition to the array $\tilde{Q}$, we



Figure 3. Network for the case of $(3, 2, 3, 2)$

obtain operators $\tilde{s}_1, \dots, \tilde{s}_{\mathfrak{b}-1}$ and π acting on $\mathcal{M}$. We emphasize that (2.7) for Q gives the operation $\tilde{\pi}$, while (2.7) for $\tilde{Q}$ gives the operation π.

The following result generalizes a result of Kajiwara, Noumi and Yamada [KNY].

Theorem 2.4. *The operators s_ℓ and π generate an action of an extended affine symmetric group $W = (\mathbb{Z}/\mathfrak{a}\mathfrak{c}\mathbb{Z}) \ltimes \widehat{\mathfrak{S}}_{\mathfrak{a}}$ on $\mathcal{M}$. Here $\mathbb{Z}/\mathfrak{a}\mathfrak{c}\mathbb{Z}$ is the cyclic group of order $\mathfrak{a}\mathfrak{c}$ and $\widehat{\mathfrak{S}}_{\mathfrak{a}}$ is the affine symmetric group (the Coxeter group) of type $\tilde{A}_{\mathfrak{a}-1}$. Similarly, the operators $\tilde{s}_\ell$ and π form an action of an extended affine symmetric group $\tilde{W} = (\mathbb{Z}/\mathfrak{b}\mathfrak{c}\mathbb{Z}) \ltimes \widehat{\mathfrak{S}}_{\mathfrak{b}}$ on $\mathcal{M}$, where $(\mathbb{Z}/\mathfrak{b}\mathfrak{c}\mathbb{Z})$ is the cyclic group of order $\mathfrak{b}\mathfrak{c}$. Furthermore, these two actions commute. Precisely, on $\mathcal{M}$ the following relations hold.*

$$\begin{aligned} & s_\ell s_{\ell+1} s_\ell = s_{\ell+1} s_\ell s_{\ell+1}, \qquad s_j s_\ell = s_\ell s_j \quad (|j-\ell|>1), \qquad s_\ell^2 = 1, \\ & \pi s_{\ell+1} = s_\ell \pi, \qquad \pi^{\mathfrak{a}\mathfrak{c}} = 1, \end{aligned}$$

$$\begin{aligned} & \tilde{s}_\ell \tilde{s}_{\ell+1} \tilde{s}_\ell = \tilde{s}_{\ell+1} \tilde{s}_\ell \tilde{s}_{\ell+1}, \qquad \tilde{s}_j \tilde{s}_\ell = \tilde{s}_\ell \tilde{s}_j \quad (|j-\ell|>1), \qquad \tilde{s}_\ell^2 = 1, \\ & \tilde{\pi} \tilde{s}_{\ell+1} = \tilde{s}_\ell \tilde{\pi}, \qquad \tilde{\pi}^{\mathfrak{b}\mathfrak{c}} = 1, \end{aligned}$$

$$s_j \tilde{s}_\ell = \tilde{s}_\ell s_j, \qquad \tilde{\pi} s_\ell = s_\ell \tilde{\pi}, \qquad \pi \tilde{s}_\ell = \tilde{s}_\ell \pi, \qquad \tilde{\pi}\pi = \pi\tilde{\pi}.$$

In the above formulae, the index of s_ℓ is taken modulo $\mathfrak{a}$, and s_0 is defined by the equation $\pi s_1 = s_0 \pi$. Similarly, the index of $\tilde{s}_\ell$ is taken modulo $\mathfrak{b}$, and $\tilde{s}_0$ is defined by the equation $\tilde{\pi}\tilde{s}_1 = \tilde{s}_0 \tilde{\pi}$.

The proof of Theorem 2.4 is delayed to §4.5.

Example 2.5. In Example 2.1 we have

$$s_1(Q)_{1,1,1} = q_{2,1,1} \frac{q_{1,1,2}q_{1,2,2}q_{1,1,1} + q_{2,2,1}q_{1,2,2}q_{1,1,1} + q_{2,2,1}q_{2,1,2}q_{1,1,1} + q_{2,2,1}q_{2,1,2}q_{2,2,2}}{q_{1,2,1}q_{1,1,2}q_{1,2,2} + q_{2,1,1}q_{1,1,2}q_{1,2,2} + q_{2,1,1}q_{2,2,1}q_{1,2,2} + q_{2,1,1}q_{2,2,1}q_{2,1,2}},$$

$$\tilde{s}_1(\tilde{Q})_{1,1,1} = q_{1,2,1} \frac{q_{1,1,1}q_{2,1,2}q_{1,1,2} + q_{1,1,1}q_{2,1,2}q_{2,2,1} + q_{1,1,1}q_{1,2,2}q_{2,2,1} + q_{2,2,2}q_{1,2,2}q_{2,2,1}}{q_{2,1,2}q_{1,1,2}q_{2,1,1} + q_{2,1,2}q_{1,1,2}q_{1,2,1} + q_{2,1,2}q_{2,2,1}q_{1,2,1} + q_{1,2,2}q_{2,2,1}q_{1,2,1}},$$

$$\tilde{\pi}(Q)_{1,1,1} = q_{1,2,1}, \qquad \pi(\tilde{Q})_{1,1,1} = q_{2,1,2}.$$

We leave it for the reader to verify that s_1 and π commute with $\tilde{s}_1$ and $\tilde{\pi}$, as an easy computational exercise.

We can present the extended affine symmetric groups W and $\tilde{W}$ as follows. The finite symmetric group $\mathfrak{S}_{\mathfrak{a}}$ acts naturally on the lattice $\mathbb{Z}^{\mathfrak{a}}$, fixing the subgroup generated by the vector $(1, 1, \dots, 1)$. Thus we have an action of $\mathfrak{S}_{\mathfrak{a}}$ on the quotient $\mathbb{Z}^{\mathfrak{a}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c})$. Then we have that $W = \mathfrak{S}_{\mathfrak{a}} \ltimes (\mathbb{Z}^{\mathfrak{a}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c}))$. The commutative normal subgroup $\mathbb{Z}^{\mathfrak{a}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c})$ is generated by e_u for $1 \le u \le \mathfrak{a}$:

$$e_u = (s_u \cdots s_{\mathfrak{a}-1})(s_{u-1} \cdots s_{\mathfrak{a}-2}) \cdots (s_1 \cdots s_{\mathfrak{a}-u})\pi^u. \tag{2.8}$$

The element e_u is identified with the vector $\mathbf{e}_u = (1, \dots, 1, 0, \dots, 0) \in \mathbb{Z}^a$ with u 1-s. Note that $e_{\mathfrak{a}} = \tilde{\pi}^{\mathfrak{a}}$ satisfies $e_{\mathfrak{a}}^{\mathfrak{c}} = 1$ agreeing with the fact that $(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c}) = 0$ in $\mathbb{Z}^{\mathfrak{a}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c})$. Similarly, we have $\tilde{W} = \mathfrak{S}_{\mathfrak{b}} \ltimes (\mathbb{Z}^{\mathfrak{b}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c}))$, where the commutative normal subgroup $\mathbb{Z}^{\mathfrak{b}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c})$ is generated by $\tilde{e}_u$ for $1 \le u \le \mathfrak{b}$:

$$\tilde{e}_u = (\tilde{s}_u \cdots \tilde{s}_{\mathfrak{b}-1})(\tilde{s}_{u-1} \cdots \tilde{s}_{\mathfrak{b}-2}) \cdots (\tilde{s}_1 \cdots \tilde{s}_{\mathfrak{b}-u})\tilde{\pi}^u. \tag{2.9}$$

Now, the operators e_u and $\tilde{e}_u$ all commute, and thus give an action of $\mathbb{Z}^{\mathfrak{a}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c}) \times \mathbb{Z}^{\mathfrak{b}}/\mathbb{Z}(\mathfrak{c}, \mathfrak{c}, \dots, \mathfrak{c})$ on $\mathcal{M}$! We will think of this as a discrete dynamical system with $\mathfrak{a} + \mathfrak{b}$ different time evolutions. We shall find a complete set of integrals of motion for this system, and study the initial value problem.

2.3. **Change of indexing.** Instead of the quadruple $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d})$, we shall also index our systems with triples (n, m, k), given by

$$n = \mathfrak{b}\mathfrak{c}, \qquad m = \mathfrak{a}, \qquad k = \mathfrak{b}\mathfrak{d}.$$

We shall also set

$$N := \gcd(n, k) = \mathfrak{b}, \qquad n' := n/N = \mathfrak{c}, \qquad k' := k/N = \mathfrak{d}, \qquad M := \gcd(n, m + k).$$

The quadruple $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d})$ can be recovered via

$$\mathfrak{a} = m, \qquad \mathfrak{b} = \gcd(n, k), \qquad \mathfrak{c} = n/\gcd(n, k), \qquad \mathfrak{d} = k/\gcd(n, k).$$

The involution $Q \longmapsto \tilde{Q}$ is associated with the following changes of indices:

$$(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) \longmapsto (\mathfrak{b}, \mathfrak{a}, \mathfrak{c}, \mathfrak{d}^{-1}),$$

$$(n, m, k) \longmapsto (mn', N, \bar{k}'m),$$

where $\bar{k}' := (k')^{-1}$ is taken modulo n'.

Through (2.2) or (2.3) we identify $q = (q_{a,b,c}) \in \mathcal{M}$ with $Q = (q_{i,j}) \in \mathrm{Mat}_{n,m}(\mathbb{C})$ or $\tilde{Q} = (\tilde{q}_{i,j}) \in \mathrm{Mat}_{mn',N}(\mathbb{C})$. Correspondingly, the network G has m vertical wires (directed upwards) which form m simple curves on the torus with the same homology class, and n horizontal wires (directed to the right) which form N simple curves on the torus with another homology class.

2.4. **Main results.** Let $\mathcal{M} \simeq \mathbb{C}^{\mathfrak{abc}} = \mathbb{C}^{mn}$ be the phase space where the ring $\mathcal{O}(\mathcal{M})$ of regular functions on $\mathcal{M}$ is generated by $q_{i,j}$ ($i \in [m], j \in [n]$). In §3, we shall define a map $\psi : \mathcal{M} \to \mathbb{C}[x, y]$. Every coefficient of $f(x, y) = \psi(q)$ is a regular function on $\mathcal{M}$. We then have (see Corollary 7.6) the following result.

Theorem. *The actions of the affine symmetric groups W and $\tilde{W}$ on $\mathcal{M}$ preserve each fiber $\psi^{-1}(f)$ for $f \in \psi(\mathcal{M})$. In particular, the coefficients of $f(x, y)$ are integrals of motion of the commuting $\mathbb{Z}^m$ and $\mathbb{Z}^N$ actions.*

In §3.2, we give a combinatorial description of every coefficient of f, as generating functions of path families on a network on the torus.

Now fix a generic $f \in \psi(\mathcal{M})$ such that the affine plane curve $\{(x, y) \mid f(x, y) = 0\} \subset \mathbb{C}^2$ is smooth (except for $(0, 0)$), and let C_f be the smooth completion of the affine curve. We shall call C_f the *spectral curve*. In §3, we define distinguished *special points*, P, A_u ($u \in [m]$) and O_u ($u \in [N]$) on C_f. We apply the results and techniques of van Moerbeke and Mumford [vMM] to establish the following; see Theorem 5.4.

Theorem. *Fix a generic $f \in \psi(\mathcal{M})$. There is an injection*

$$\phi : \psi^{-1}(f) \hookrightarrow \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A,$$

where

1. *g is the genus of C_f, and $\mathrm{Pic}^g(C_f)$ is the Picard group of C_f, of degree g;*
2. *$\mathcal{S}_f \simeq (\mathbb{C}^*)^{M-1} \subset (\mathbb{C}^*)^M$;*
3. *R_O is a finite set of cardinality $N!$, identified with the $N!$ orderings of $\{O_1, O_2, \ldots, O_N\}$; and*
4. *R_A is a finite set of cardinality $m!$, identified with the $m!$ orderings of $\{A_1, A_2, \ldots, A_m\}$.*

In fact, our Theorem 5.4 identifies the image of ϕ.

Define the divisors $\mathcal{A}_u := uP - \sum_{i=1}^{u} A_i$ and $\mathcal{O}_u := uP - \sum_{j=N-u+1}^{N} O_j$. The commuting time evolutions e_u (2.8) and $\tilde{e}_u$ (2.9) can be described as follows (Theorem 7.2 and Theorem 7.3).

Theorem. *Suppose $\phi(q) = ([\mathcal{D}], (c_1, \ldots, c_M), O, A)$. Then we have*

$$\phi(e_u(q)) = ([\mathcal{D} - \mathcal{A}_u], (c_{u+1}, \ldots, c_M, c_1, \ldots, c_u), O, A) \quad \text{for } u = 1, \ldots, m,$$
$$\phi(\tilde{e}_u(q)) = ([\mathcal{D} + \mathcal{O}_u], (c_{M-u+1}, \ldots, c_M, c_1, \ldots, c_{M-u}), O, A) \quad \text{for } u = 1, \ldots, N.$$

Furthermore, the finite symmetric subgroups $\mathfrak{S}_N \subset \tilde{W}$ and $\mathfrak{S}_m \subset W$ act naturally on R_O and R_A respectively and do not affect the rest of the spectral data.

In other words, the time evolutions e_u and $\tilde{e}_u$ are linearized on $\mathrm{Pic}^g(C_f)$. Our approach to Theorem 7.3 is similar to that of Iwao [I08].

When $N = 1$, we give in Theorem 8.5 an explicit formula for the inverse to the map ϕ, which also explicitly solves the initial value problem for our dynamics. For $t \in \mathbb{Z}^m$, let $(q_{j,i}^t)$ denote the point in the phase space after time evolution in the direction t.

Theorem. *When $N = 1$, we have a formula*

$$q_{j,i}^t = C \, \frac{\theta_i^{t+\mathbf{e}_j}(P) \, \theta_{i-1}^{t+\mathbf{e}_{j-1}}(P)}{\theta_i^{t+\mathbf{e}_{j-1}}(P) \, \theta_{i-1}^{t+\mathbf{e}_j}(P)},$$

where $\theta_i^t(P)$ *is a particular value of the Riemann theta function, and* C *is a constant depending only on* $(c_1, \dots, c_M)$*,* O*, and* A*.*

2.5. **Examples.**

2.5.1. *Discrete Toda lattice.* Let $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (2, 1, n, n-1)$ (i.e. $(n, m, k) = (n, 2, n-1)$). We set $q_{i,j} = q_{a,1,c}$ for $i = a,\ j = c$, and regard $q := (q_{i,j})_{i\in[2],j\in[n]}$ as a coordinate of $\mathcal{M} \simeq \mathbb{C}^{2n}$. The resulting $\mathbb{Z}^2$-action on $\mathcal{M}$ is generated by e_1 and e_2, where e_2 simply acts on $\mathcal{M}$ as $e_2(q_{i,j}) = q_{i,j+1}$. As for e_1, when we define $q^t := e_1^t(q)$ for $q \in \mathcal{M}$, the action of e_1 is rewritten as a system of difference equations:

$$\begin{cases} q_{1,j}^{t+1} q_{2,j}^{t+1} = q_{2,j}^t q_{1,j+1}^t, \\ q_{1,j+1}^{t+1} + q_{2,j}^{t+1} = q_{2,j+1}^t + q_{1,j+1}^t. \end{cases} \tag{2.10}$$

This is the discretization of n-periodic Toda lattice equation studied in [HTI]. Actually, we recover the original Toda lattice equation $\frac{d^2}{dt^2}x_j = \mathrm{e}^{x_{j+1}-x_j} - \mathrm{e}^{x_j - x_{j-1}}$ by setting $q_{1,j}^t = 1 + \delta \frac{d}{dt} x_j$ and $q_{2,j}^t = \delta^2 \mathrm{e}^{x_{j+1}-x_j}$, and taking the limit $\delta \to 0$. Here we set $q_{*,j}^t = q_{*,j}(\delta t)$ for $* = 1, 2$ and $x_j = x_j(t)$.

A simple generalization of the discrete Toda lattice is the case of $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (m, 1, n, n-1)$. Its initial value problem was studied by Iwao [I08, I10] by applying [vMM]. He also studied the similar problem in the case of $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (m, 1, n, 0)$ in [I09].

2.5.2. *Tropicalization and tableaux.* Let T_1 and T_2 be two semistandard tableaux of rectangular shapes. Define $T_1 \otimes T_2$ to be the concatenation of the two into a skew tableaux by placing T_2 North-East of T_1. Let $\mathrm{jdt}(T_1 \otimes T_2)$ denote the straight shape tableau obtained by performing jeu-de-taquin on $T_1 \otimes T_2$. The following result is well-known.

Lemma 2.6. *There is a unique pair of rectangular semistandard tableaux* T_1' *and* T_2' *such that* T_i' *has the same shape as* T_i *(for* $i = 1, 2$*), and* $\mathrm{jdt}(T_1 \otimes T_2) = \mathrm{jdt}(T_2' \otimes T_1')$.

Example 2.7. Suppose

$$T_1 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 3 \\ \hline \end{array} \qquad \text{and} \qquad T_2 = \begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline \end{array}.$$

Then one has

$$T_2' = \begin{array}{|c|c|c|} \hline 1 & 3 & 3 \\ \hline \end{array} \qquad \text{and} \qquad T_1' = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 2 & 3 \\ \hline \end{array}.$$

The transformation $R(T_1, T_2) = (T_2', T_1')$ is called the *combinatorial R-matrix.* It appears in the theory of crystal graphs as the isomorphism map between tensor products of Kirillov-Reshetikhin crystals. The following property is well-known.

Theorem 2.8. *The combinatorial R-matrix is an involution. Furthermore, it satisfies the braid relation:*

$$(R \otimes Id)(Id \otimes R)(R \otimes Id) = (Id \otimes R)(R \otimes Id)(Id \otimes R).$$

Now, let $\{q_{a,b,c}\}_{a\in[\mathfrak{a}],b\in[\mathfrak{b}],c\in[\mathfrak{c}]}$ be an array of nonnegative integers. Define $Q = (q_{i,j})$ and $\tilde{Q} = (\tilde{q}_{i,j})$ as before:

$$q_{i,j} = q_{a,b,c} \text{ for } i = a,\ j = b + \mathfrak{b}(c-1)$$

and

$$\tilde{q}_{ij} = q_{a,b,c} \text{ for } i = \mathfrak{b} + 1 - b,\ j = \mathfrak{a}\mathfrak{d}^{-1}c - a + 1.$$

We create single-row tableaux from Q and $\tilde{Q}$ as follows. For each $i \in [\mathfrak{a}]$, let T_i be the single-row tableau with $q_{i,j}$ j-s. For each $i \in [\mathfrak{b}]$, let $\tilde{T}_i$ be the single-row tableau with $\tilde{q}_{i,j}$ j-s. We view $T = T_1 \otimes \ldots \otimes T_{\mathfrak{a}}$ and $\tilde{T} = \tilde{T}_1 \otimes \ldots \otimes \tilde{T}_{\mathfrak{b}}$ as tropical analogues of Q and $\tilde{Q}$.

Let us define an action of W on $q_{a,b,c}$ as follows. For $T = T_1 \otimes \ldots \otimes T_{\mathfrak{a}}$ and $1 \leq \ell \leq \mathfrak{a}-1$ let

$$s_\ell(T) = T_1 \otimes \ldots \otimes T'_{\ell+1} \otimes T'_\ell \otimes \ldots \otimes T_{\mathfrak{a}}.$$

That is, we apply the combinatorial R-matrix to the ℓ-th and $(\ell+1)$-st factors. In addition, let $\tilde{\pi}$ act on T by applying Schutzenberger's *promotion* operator [Sch] to each factor T_i. Similarly, we define $\tilde{s}_\ell$ and π acting on $\tilde{T}$.

Theorem 2.9. *The operators s_ℓ and π form an action of the extended affine symmetric group W on $q_{a,b,c}$. Similarly, operators $\tilde{s}_\ell$ and $\tilde{\pi}$ form an action of the extended affine symmetric group $\tilde{W}$ on $q_{a,b,c}$. These two actions commute. These two operations are the tropicalizations of the rational actions of Theorem 2.4.*

Here tropicalization refers to the formal operation of substitution

$$\mathbb{C} \mapsto \mathbb{Z}, \qquad \times \mapsto +, \qquad \div \mapsto -, \qquad + \mapsto \min .$$

Under this substitution, the geometric R-matrix becomes a piecewise-linear involution of $\mathbb{Z}^n \times \mathbb{Z}^n$, which is the combinatorial R-matrix. In other words, the W dynamics we are considering in this paper is a birational lift of the dynamics of repeated application of the combinatorial R-matrix on a sequence of $\mathfrak{a}$ single-row tableaux, arranged in a circle.

Example 2.10. Let $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (2, 2, 2, 1)$ as in Example 2.1. Take $T = 22222344 \otimes 1234$. Then $\tilde{T} = 122222344 \otimes 134$. We have

$$s_1(T) = 2234 \otimes 12222344, s_1(\tilde{T}) = 111122334 \otimes 134;$$

$$\tilde{\pi}(T) = 11111233 \otimes 1234, \tilde{\pi}(\tilde{T}) = 123 \otimes 122222344;$$

$$\tilde{s}_1(T) = 11112334 \otimes 1233, \tilde{s}_1(\tilde{T}) = 124 \otimes 122223344;$$

$$\pi(T) = 1234 \otimes 22222344, \pi(\tilde{T}) = 111112334 \otimes 234.$$

Remark 2.11. The corresponding commuting crystal actions in the case $\mathfrak{c} = 1$ were considered by Lascoux in [Las] and by Berenstein-Kazhdan in [BK].

2.5.3. *Box-ball systems.* The box-ball system is an integrable cellular automaton introduced by Takahashi and Satsuma [TS]. It is described by an algorithm to move finitely many balls in an infinite number of boxes aligned on a line, where a consecutive array of occupied boxes is regarded as a *soliton.* This system is related to both of the previous two examples; the global movements of solitons are equivalent to the tropicalization of the discrete Toda lattice (2.10). The symmetry of the system is explained by the crystal base theory, where the dynamics of bolls is induced by the action of the combinatorial R-matrix.

See [IKT] for a comprehensive review of the combinatorial and tropical aspects of the box-ball system.

3. Lax matrix and spectral curve

3.1. **Lax matrix.** Fix integers n, m, k such that $n \geq 2$, $m \geq 1$ and $1 \leq k \leq n$. From now on we shall mainly use $q := (q_{ij})_{i \in [m], j \in [n]}$ as a coordinate of the phase space $\mathcal{M}$. We identify $q \in \mathcal{M}$ with an m-tuple of n by n matrices $Q := (Q_i(x))_{i=1,\ldots,m}$ with a spectral parameter x, where

$$Q_i(x) := \begin{pmatrix} q_{i1} & 0 & 0 & x \\ 1 & q_{i2} & 0 & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & 0 & 1 & q_{in} \end{pmatrix}. \tag{3.1}$$

Let $\mathcal{L}$ be the set of n by ∞ scalar matrices $A := (a_{ij})_{1 \leq i \leq n,\, j \in \mathbb{Z}}$ satisfying the following conditions:

$$a_{ij} = \begin{cases} 1 & j - i = -m - k \\ a_{ij} \in \mathbb{C} & -m - k + 1 \leq j - i \leq -k \\ 0 & \text{otherwise.} \end{cases} \tag{3.2}$$

In particular, A has finitely many nonzero entries. For $A \in \mathcal{L}$, we define an n by n matrix $L(A; x) = (l(x)_{ij})_{1 \leq i,j \leq n}$ by

$$l(x)_{ij} = \sum_{\ell \in \mathbb{Z}} a_{i, j - \ell n}\, x^{\ell}. \tag{3.3}$$

We may identify $\mathcal{L}$ with $\mathbb{C}^{mn}$, and identify $A \in \mathcal{L}$ with $L(A; x)$.

Now define a map $\alpha : \mathcal{M} \to \mathcal{L}$ by

$$\alpha : Q = (Q_1, Q_2, \ldots, Q_m) \longmapsto L(x) := Q_1(x) Q_2(x) \cdots Q_m(x) P(x)^k,$$

where

$$P(x) := \begin{pmatrix} 0 & 0 & 0 & x \\ 1 & 0 & 0 & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}. \tag{3.4}$$

We call $L(x)$ the *Lax matrix*. Our approach to the study of $L(x)$ is close to that of van Moerbeke and Mumford [vMM].

We give a combinatorial description of the Lax matrix by using *highway paths* on the network G. We introduce the "x-line" as illustrated in Figure 4, where the top k right ends of horizontal wires cross the x-line (the top edge) and come out on the other side. There are n sources labeled $1, 2, \ldots, n$ on the left and n sinks labeled $1, 2, \ldots, n$ on the right. Each sink j is connected to the source $j + k \mod n$ by a wire, as illustrated in the figure. There are mn intersection points between the vertical wires and the horizontal wires, which we call the (i, j)-*crossroads*, where $i = 1, 2, \ldots, m$ indexes the vertical wire, and $j = 1, 2, \ldots, n$ indexes the horizontal wire. Let us denote by G' the *cylindrical network* obtained by gluing only the upper and lower edges of Figure 4. In the torus network G, source i and sink i are the same point. In the cylindrical network G', source i and sink i are distinct points.

Let us introduce the notion of highway paths following [LP13]. A *highway path* p is a directed path in the network G (or G') with the following property: at any of the



FIGURE 4. The toric network G, with a coil and a snake path shown.

crossroads, if the path is traveling upwards, it must turn right. We shall only consider highway paths that start at one of the sources $1, 2, \dots, n$ and end at one of the sinks $1, 2, \dots, n$. The weight $\mathrm{wt}(p)$ of a highway path p is defined as follows. Every time p passes the x-line it picks up the weight x. Every time p goes through the (i, j)-crossroad, it picks up the weight q_{ij} or 1, according to Figure 5. The weight $\mathrm{wt}(p)$ is the product of all these weights. The condition that a highway path must turn right when travelling up into a crossroad is indicated in the Figure 5: we can think of such a turn as giving weight 0. Finally, a highway path p may use no edges. In this case, we consider the path p to start at some source i, and end at the sink i. We declare such paths to be *abrupt*, and have weight $\mathrm{wt}(p) = -y$.



FIGURE 5. Highway paths

The following lemma gives a highway-path description of the Lax matrix, which is a variant of the results of [LP13]. It follows directly from the definitions.

Lemma 3.1. *Let $L(x) = \alpha(q)$ for $q \in \mathcal{M}$. For $1 \le i, j \le n$, we have*

$$(i,j)\text{-th entry of } L(x) - y = \sum_p \mathrm{wt}(p),$$

where the summation is over highway paths in G' from source i to sink j.

3.2. **Spectral curve and Newton polygon.** Define a map $\psi : \mathcal{M} \to \mathbb{C}[x, y]$ as the composition of $\alpha : \mathcal{M} \to \mathcal{L}$ and the map $\beta : \mathcal{L} \to \mathbb{C}[x, y]$,

$$Q = (Q_1, Q_2, \dots, Q_m) \xmapsto{\alpha} L(x) = Q_1(x)Q_2(x) \cdots Q_m(x)P(x)^k \xmapsto{\beta} \det(L(x) - y).$$

Consequently, for $Q = (Q_1, \dots, Q_m) \in \mathcal{M}$ we have an affine plane curve $C'_{\psi(Q)}$ in $\mathbb{C}^2$, given by the zeros of $\psi(Q)$. We call this curve the *spectral curve*.

Each term of $\psi(Q)$ corresponds to the weight of specific highway paths as follows. We say that a pair of paths is *noncrossing* if no edge is used twice, and that a family of paths is noncrossing if every pair of paths is noncrossing. Suppose $\mathbf{p} = \{p_1, p_2, \ldots, p_n\}$ is an unordered noncrossing family of n paths in G' using all the sources and all the sinks. The non-abrupt paths in $\mathbf{p}$ induce a bijection of a subset $S \subset [n]$ with itself. We let $\mathrm{sign}(\mathbf{p})$ denote the sign of this permutation. The following theorem is a reformulation [MIT]. In our language, the proof is very similar to [LP12a, Theorem 3.5].

Theorem 3.2. *We have*

$$f(x,y) = \det(L(x) - y) = \sum_{\mathbf{p}=\{p_1,p_2,\ldots,p_n\}} \mathrm{sign}(\mathbf{p})\mathrm{wt}(p_1)\mathrm{wt}(p_2)\cdots\mathrm{wt}(p_n),$$

where the summation is over noncrossing (unordered) families of n paths in G' using all the sources and all the sinks. In other words, the coefficient $x^a y^b$ in $f(x,y) = \det(L(x) - y)$ counts (with weights) families of n paths that

- *do not cross each other;*
- *cross the x-line exactly a times;*
- *contain exactly b abrupt paths and $n - b$ non-abrupt paths.*

The overall resulting sign of the monomial $x^a y^b$ is $(-1)^{(n-b-1)a+b}$.

For $f(x,y) = \sum_{i,j} a_{i,j} x^i y^j \in \mathbb{C}[x,y]$, we write $N(f) \subset \mathbb{R}^2$ for the Newton polygon of f. This is defined to be the convex hull of the points $\{(i,j) \mid a_{ij} \neq 0\}$. It is important for us to identify the *lower hull* and *upper hull* of $N(f)$. The former (resp. latter) is the set of edges of $N(f)$ such that the points directly below (resp. above) these edges do not belong to $N(f)$. We exclude vertical or horizontal edges from the lower and upper hull.

Proposition 3.3. *For generic $Q = (Q_1, \ldots, Q_m) \in \mathcal{M}$, the Newton polygon $N(\psi(Q))$ is the triangle with vertices $(0,n)$, $(k,0)$ and $(k+m,0)$. In $N(\psi(Q))$, the lower hull (resp. upper hull) consists of one edge with vertices $(k,0)$ and $(0,n)$ (resp. $(k+m,0)$ to $(0,n)$).*

See § 4.1 for the proof.

3.3. **Special points on the spectral curve.** For $f(x,y) \in \mathbb{C}[x,y]$ an irreducible polynomial, let $C'_f = \{(x,y) \mid f(x,y) = 0\} \subset \mathbb{C}^2$ be the corresponding plane curve, and let $\overline{C'_f} \subset \mathbb{P}^2(\mathbb{C})$ denote its closure. Let C_f be the normalization of $\overline{C'_f}$, with a map $C_f \to \overline{C'_f}$ that is a resolution of singularities. We declare points on C_f to be *special* if either (1) either x or y is 0, or (2) the point does not lie over C'_f (that is, x or y is ∞).

For $f(x,y) \in \psi(\mathcal{M})$, define a polynomial $g_f(x,y)$ (resp. $h_f(x,y)$) by $f(x,y) = g_f(x,y) + \text{other terms}$ (resp. $f(x,y) = h_f(x,y) + \text{other terms}$), where $g_f(x,y)$ (resp. $h_f(x,y)$) consists of those monomials lying on the lower hull (resp. the upper hull) of $N(f)$. For this $f(x,y)$ we also define

$$f_c := \sum_{(j,i)\in L_c} f_{i,j}, \tag{3.5}$$

where

$$L_c = \{(j,i) \mid (m+k)i + nj = n(m+k) - M\}.$$

We define $\sigma_r \in \mathcal{O}(\mathcal{M})$ by

$$\sigma_r(q) = \prod_{i=1}^{m} \prod_{j=0}^{n'-1} q_{i,r+jN}, \tag{3.6}$$

for $1 \leq r \leq N$, and define $\epsilon_r \in \mathcal{O}(\mathcal{M})$ by

$$\epsilon_r(q) := \prod_{j=1}^{n} q_{rj} \tag{3.7}$$

for $1 \leq r \leq m$.

Remark 3.4. Despite the seeming dissimilarity, $\sigma_r(q)$ and $\epsilon_r(q)$ have the same nature. Indeed, visualize variables q_{ij} as associated to crossings of two families of parallel wires on a torus, as it was done in Section 2. Then $\sigma_r(q)$ is the product of parameters on the r-th horizontal wire (out of N), while $\epsilon_r(q)$ is the product of parameters on the r-th vertical wire (out of m). In particular, the symmetry between Q and $\tilde{Q}$ from Section 2 switches the $\sigma_r(q)$ and the $\epsilon_r(q)$ into each other.

Lemma 3.5. *For $f(x,y) \in \psi(\mathcal{M})$, we have*

$$g_f(x,y) = \prod_{r=1}^{N}((-y)^{n'} + \sigma_r x^{k'}), \tag{3.8}$$

$$h_f(x,y) = ((-y)^{\frac{n}{M}} + x^{\frac{k+m}{M}})^M. \tag{3.9}$$

See § 4.2 for the proof.

In the following, for $f(x,y) = \psi(q),\ q \in \mathcal{M}$ we study the special points of C_f related to polynomials $g_f(x,y)$, $h_f(x,y)$ or $f(x,0)$.

We have $f(x,0) = \det P(x)^k \prod_{i=1}^m \det Q_i(x)$, where $\det Q_i(x) = \prod_{j=1}^n q_{ij} + (-1)^{n-1}x = \epsilon_i(q) + (-1)^{n-1}x$. Thus the roots of $f(x,0) = 0$ are exactly the $(-1)^n \epsilon_r(q)$, where $\epsilon_r(q)$ is defined by (3.7). It is also clear that $\epsilon_1, \ldots, \epsilon_m$ depend only on $f(x,y)$. The following lemma is obtained immediately.

Lemma 3.6. *Suppose that $\epsilon_1, \ldots, \epsilon_m$ are distinct and nonzero. Then there are m special points $A_i = ((-1)^n \epsilon_i(q), 0)$ on C_f with $y = 0$ and x is nonzero.*

The point $(0,0)$ on C'_f is usually singular, whenever $k \geq 2$.

Lemma 3.7. *Suppose $\sigma_1, \ldots, \sigma_N$ are distinct and nonzero. Then there are N points of C_f lying over $(0,0) \in C'_f$.*

Proof. By (3.8), the meromorphic function $y^{n'}/x^{k'}$ takes the N distinct values $(-1)^{n'+1}\sigma_r$ for $r = 1, 2, \ldots, N$ as $(x,y) \to (0,0)$. So there are at least N points on C_f. On the other hand, looking at $N(f)$ we see that analytically near $(0,0)$, the polynomial f can factor into at most N pieces. □

Let $O_1, O_2, \ldots, O_N$ denote the the special points of Lemma 3.7. Near O_r there is a local coordinate u such that

$$(x,y) \sim (u^{n'}, -(-\sigma_r)^{1/n'} u^{k'}).$$

It turns out that $C'_f \subset \mathbb{P}^2(\mathbb{C})$ has only one point at ∞. Due to the polynomial $h_f(x,y)$, in homogeneous coordinates, this point is

$$P' = \begin{cases} [1:0:0] & \text{if } n > k+m, \\ [1:1:0] & \text{if } n = k+m, \\ [0:1:0] & \text{if } n < k+m. \end{cases}$$

To compute this, we first homogenize $f(x,y)$ to get $F(x,y,z) = z^d f(x/z, y/z)$where $d := \max(n, m+k)$. Then we solve $F(x,y,0) = 0$. Recall that f_c was defined in (3.5).

Lemma 3.8. *Suppose that $f_c \neq 0$. Then there is a unique point $P \in C_f$ lying over P'.*

See § 4.3 for the proof.

3.4. **A good condition for the spectral curve.**

Let $V_{n,m,k}$ be the subspace of $\mathbb{C}[x,y]$ given by

$$V_{n,m,k} = \left\{ ((-y)^{\frac{n}{M}} + x^{\frac{k+m}{M}})^M + \sum_{i=0}^{n-1} y^i f_i(x) \;\middle|\; f_i(x) = \sum_{(j,i)\in L_{n,m,k}} f_{i,j} x^j \in \mathbb{C}[x] \right\}, \quad (3.10)$$

where we write $L_{n,m,k}$ for the set of lattice points in the convex hull of $\{(k,0),(0,n),(k+m,0)\}$ but not on the upper hull. By Proposition 3.3 and Lemma 3.5, we have that $\psi(\mathcal{M}) \subset V_{n,m,k}$.

Definition 3.9. Define the subset $\mathcal{V} \subset V_{n,m,k}$ as the set of $f(x,y) \in V_{n,m,k}$, satisfying the conditions:

1. $f(x,y)$ is irreducible;
2. C'_f is smooth;
3. $f_c \neq 0$;

and such that the special points on C_f consist exactly of:

4. m distinct points $A_1, A_2, \ldots, A_m$ where $A_r := ((-1)^n \epsilon_i, 0)$ and $\epsilon_i \neq 0$;
5. N distinct points $O_1, O_2, \ldots, O_N$ lying over $(0,0)$, where near O_r there is a local coordinate u such that
$$(x,y) \sim (u^{n'}, -(-\sigma_r)^{1/n'} u^{k'}) \quad (3.11)$$
and $\sigma_r \neq 0$;
6. a single point P lying over the line at infinity $\mathbb{P}^2(\mathbb{C}) \setminus \mathbb{C}^2$.

For $f \in \mathcal{V}$, the genus g of C_f is given by

$$g = \frac{1}{2}\left((n-1)m - M - N + 2\right). \quad (3.12)$$

Indeed, it follows from Pick's formula and Proposition 3.3 that the number of interior lattice points of $N(f)$ is equal to the right hand side. This formula for the genus then follows from [Kho, Corollary on p.6]. Alternatively, the genus can be computed using the Riemann-Hurwitz formula, as in [vMM].

Proposition 3.10. *For $f \in \mathcal{V}$, we have that $\beta^{-1}(f) \neq \emptyset$. Moreover, the set $\mathring{\mathcal{V}} := \mathcal{V} \cap \psi(\mathcal{M})$ is a Zariski-dense subset of $\psi(\mathcal{M})$.*

We give the proof in § 4.4. Informally, the last statement of Proposition 3.10 states that most curves in $\psi(\mathcal{M})$ satisfy the "niceness" conditions listed in Definition 3.9.

From Definition 3.9, it follows that for $f \in \mathcal{V}$, the meromorphic functions x and y on C_f satisfy

$$(x) = n' \sum_{i=1}^{N} O_i - nP, \qquad (y) = k' \sum_{i=1}^{N} O_i + \sum_{i=1}^{m} A_i - (m+k)P. \quad (3.13)$$

Remark 3.11. The condition that $A_1, \ldots, A_m$ (resp. $O_1, O_2, \ldots, O_N$) are distinct imply that the quantities $\epsilon_1, \ldots, \epsilon_m$ (resp. $\sigma_1, \ldots, \sigma_N$) are distinct. Many of our main results still apply after a modification even when this condition does not hold.

4. Proofs from Sections 2 and 3

4.1. **Proof of Proposition 3.3.** We use the interpretation of $f(x, y)$ given by Theorem 3.2. Let $\mathbf{p} = (p_1, \ldots, p_n)$ be a family of noncrossing highway paths in G', as in Theorem 3.2, and let $\mathrm{wt}(\mathbf{p}) = \mathrm{wt}(p_1) \cdots \mathrm{wt}(p_n)$.

There are $k + m$ opportunities for a highway path in $\mathbf{p}$ to pick up the weight x: from each of the m vertical wires and from the k horizontal wires crossing x-line. For a fixed power y^b (where $b = 0, 1, \ldots, n$), or equivalently a fixed number of abrupt paths, we will bound the maximal and the minimal possible value of the exponent a of x.



Figure 6. The red path crosses x-line less than the green one.

The first key observation is that we can consider $\mathbf{p}$ to be a family of noncrossing closed cycles on the toric network G. An abrupt path p is simply the "cycle" that starts and ends at the vertex i (= source i and sink i identified) in G and does not move.

Lift such a highway cycle C to the universal cover, as shown in Figure 6. We obtain a path that starts and ends at two vertices labeled with the same integer $i \in \{1, 2, \ldots, n\}$, but ℓ periods (of m vertical lines each) to the right of the original source. More precisely, if C is obtained by gluing paths $p_{i_1}, \ldots, p_{i_\ell}$ in G', then C has length ℓ. The x-line now has a staircase-like shape as shown in Figure 6. We claim that if C has length ℓ, then it crosses the x-line at least $\ell k/n$ times and at most $\ell(k + m)/n$ times.

To see this, observe that if we lift the ending vertex several periods (consisting of n horizontal wires) up, we preserve the length and we increase the number of crossings with the x-line. Similarly, if we lower the ending vertex several periods down, we preserve the length and we decrease the number of crossings of x-line. Thus, the smallest and the largest ratios of the quantity (crosses of x-line/length) is achieved for the lowest and the highest possible highway paths, shown in Figure 6 in purple and green. The former is the horizontal path, while the latter is an alternating right-up staircase path. For these paths, the ratios are exactly k/n and $(k+m)/n$.

4.2. **Proof of Lemma 3.5.** One of the consequences of the proof of Proposition 3.3 is that the monomials for which the lower bound k/n of the ratio is reached are the ones coming from horizontal highway paths on the universal cover. Let us call each such closed cycle a *coil.* For example, the purple line in Figure 6 represents a coil passing through source i, as well as passing through the n' other vertices between 1 and n that have residue i modulo $N = \gcd(k,n)$. Thus the terms of $g(x,y)$ are formed in the following way: for each of the N coils we decide whether to include it into our family of paths, or to make all paths starting at its sources abrupt. The second choice corresponds to the contribution $(-y)^{n'}$. The first choice gives $(\prod_{i=1}^{m}\prod_{j=1}^{n'} q_{i,r+jN})x^{k'}$, which is the weight of that coil. Thus, the r-th coil contributes the factor of $\left((\prod_{i=1}^{m}\prod_{j=1}^{n'} q_{i,r+jN})x^{k'} + (-y)^{n'}\right)$, and (3.8) follows.

Similarly, the monomials for which the upper bound $(k+m)/n$ of the ratio is reached are the ones coming from right-up staircase paths on the universal cover. Let us call each such closed cycle a *snake path.* For example, the green line in Figure 6 represents a snake path passing through source i, as well as through all the n/M other vertices between 1 and n that have residue i modulo $M = \gcd(m+k,n)$. Thus the part contributing to the upper hull of $f(x,y)$ is a product of factors $x^{(m+k)/M} + (-y)^{n/M}$, where the term $(-y)^{n/M}$ corresponds to choosing to have abrupt paths, while $x^{(m+k)/M}$ corresponds to choosing to have the snake path. Thus we obtain (3.9).

4.3. **Proof of Lemma 3.8.** We analyze the singularity at $P' \in \mathbb{P}^2(\mathbb{C})$ on the chart $y \neq 0$, that is, $\{[x:1:z] \mid x,z \in \mathbb{C}\}$. Let $\tilde{C}_{\overline{f}}$ be the affine curve given by $\overline{f}(x,z) := F(x,1,z) = 0$.

(i) $n < k+m$: using (3.9) we can write $\overline{f}(x,z)$ as

$$\overline{f}(x,z) = ((-1)^{\frac{n}{M}} z^{\frac{k+m-n}{M}} + x^{\frac{k+m}{M}})^M + \sum_{j=1}^{\max(m,m+k-n)} z^j \overline{f}_j(x)$$

where $\deg_x \overline{f}_j(x) \geq 1$. Then, when $m+k-n = 1$, $\tilde{C}_{\overline{f}}$ is smooth at $(0,0)$ since $\partial\overline{f}/\partial z|_{(0,0)} = (-1)^{n/M} \neq 0$. When $m+k-n \geq 2$, the point $(0,0)$ is singular. Define $K := \mathbb{C}(x)[z]/(\overline{f}(x,z))$. We consider all valuations $v : K \twoheadrightarrow \mathbb{Z}$ on K satisfying that $v(x) > 0$ and $v(z) > 0$. We shall show that such valuations are unique, from which the uniqueness of P follows. Define $\overline{h}_f(x,z) := ((-1)^{\frac{n}{M}} z^{\frac{k+m-n}{M}} + x^{\frac{k+m}{M}})^M$ which is the part of $\overline{f}(x,z)$ consisting of monomials lying on the lower hull of $N(\overline{f})$, corresponding to the upper hull of $N(f)$. Then we see that $v(\overline{f}(x,z)) = v(\overline{h}_f(x,z)) = M \cdot v(x^{\frac{k+m}{M}} + (-1)^{\frac{n}{M}} z^{\frac{k+m-n}{M}})$. To be consistent with $v(\overline{f}(x,z)) = v(0) = \infty$, the only choice we have is $v(x) = \frac{k+m-n}{M}$ and $v(z) = \frac{k+m}{M}$.

(ii) $n = k+m$: we can write $\overline{f}(x,z)$ as

$$\overline{f}(x,z) = (x-1)^n + \sum_{k=1}^{m} z^k \overline{f}_k(x-1)$$

where $\overline{f}_k(x-1) \in \mathbb{C}[x-1]$. Thus P' is smooth point on $C_{\overline{f}}$ if $\overline{f}_1(0) \neq 0$. Looking at the Newton polygon $N(f)$ and (3.5), it follows that $\overline{f}_1(0) = \sum_{i=0}^{n-1} f_{i,n-1-i} = f_c \neq 0$.

(iii) $n > k+m$: this case is nearly the same as the case $n < k+m$.

4.4. **Proof of Proposition 3.10.** The first statement follows from Theorem 5.3 in the next section. In the following, we prove the second statement.

We first show that $\mathcal{V}$ contains a Zariski-dense and open subset of $V_{n,m,k}$. A Zariski-open subset of $V_{n,m,k}$ consists of $f(x,y)$ with Newton polygon $N(f)$ given by Proposition 3.3. This polygon is not a non-trivial Minkowski sum of two other polygons, so $f(x,y)$ is irreducible. Similarly the conditions that C'_f is smooth and $f_c \neq 0$ are Zariski-open conditions on $V_{n,m,k}$. By Lemma 3.8, $f_c \neq 0$ implies that (6) in Definition 3.9 holds. The calculations in the proofs of Lemmas 3.6 and 3.7 then imply that $\mathcal{V}$ contains a Zariski-open and dense subset of $V_{n,m,k}$.

It thus suffices to show that $\psi(\mathcal{M})$ contains a Zariski-dense subset of $V_{n,m,k}$. We shall use the following result.

Lemma 4.1. *The set $\alpha(\mathcal{M})$ is Zariski-dense in $\mathcal{L}$.*

Proof. This follows from [LP11, Proof of Theorem 4.1] where it is shown that the map $\alpha : \mathcal{M} \to \mathcal{L}$ is generically a $m!$ to 1 map between two spaces of dimension mn. ☐

By the first statement of Proposition 3.10, we have $\mathcal{V} \subset \beta(\mathcal{L})$. By Lemma 4.1, $\psi(\mathcal{M}) = \beta(\alpha(\mathcal{M}))$ contains a Zariski-dense subset of $\mathcal{V}$. It follows that $\psi(\mathcal{M})$ is Zariski-dense in $V_{n,m,k}$. This completes the proof of the second statement of Proposition 3.10.

Remark 4.2. The nonconstant coefficients $f_{i,j}$ of $f(x,y)$ can be pulled back to functions on $\mathcal{M}$ or $\mathcal{L}$. A consequence of our proof is that $\psi(\mathcal{M})$ is Zariski-dense in $V_{n,m,k}$, and therefore the functions $f_{i,j}$ are algebraically independent on $\mathcal{M}$ or $\mathcal{L}$. It would be interesting to obtain a direct proof of this.

4.5. **Proof of Theorem 2.4.** We employ the technique first introduced in [LP13].



Figure 7. Crossing merging and crossing removal moves with vertex weights shown.



Figure 8. Yang-Baxter move with transformation of vertex weights shown.

Specifically, we realize the geometric R-matrix transformation, dubbed the *whirl move* in [LP13], as a sequence of local transformations on our toric network. The transformations we shall employ are of three kinds: crossing merging/unmerging, crossing creation/removal, shown in Figure 7, and Yang-Baxter move shown in Figure 8.

The whirl move R occurs between two parallel wires adjacent to each other and wrapping around a local part of the surface that is a cylinder. The way it is realized as a sequence of local moves is illustrated in Figure 9. First a crossing is created with weight 0, and split into two crossings of weight p and $-p$. One of them is pushed through the wires crossing our two distinguished wires, until it comes out on the other side. As it is proven in [LP13, Theorem 6.2], there is at most one non-zero value of p for which the end result is again a pair of crossings of weights p and $-p$, and thus those two can be canceled out. The resulting action on the weights along two parallel wires is exactly the whirl move R, and does not depend on where the original auxiliary crossing was created. The value of p does depend on the location j where the new crossings are created, and is given by



FIGURE 9. For a unique choice of the weight p, the weight that comes out on the other side after passing through all horizontal wires is also p; the resulting transformation of x and y is exactly the whirl move.

$$p = \frac{\prod y_i - \prod x_i}{E(\boldsymbol{x}^{(j)}, \boldsymbol{y}^{(j)})}.$$

Here the variables x_i and y_i are weights along the two wires as in Figure 9, and $E(\boldsymbol{x}^{(j)}, \boldsymbol{y}^{(j)})$ is the energy of the cyclically shifted vectors $\boldsymbol{x}^{(j)} = (x_{j+1}, x_{j+2}, \ldots, x_j)$ and $\boldsymbol{y}^{(j)} = (y_{j+1}, y_{j+2}, \ldots, y_j)$ as introduced in § 2.1.

Now, the fact that R satisfies the braid move, that is, $s_\ell s_{\ell+1} s_\ell = s_{\ell+1} s_\ell s_{\ell+1}$ and $\tilde{s}_\ell \tilde{s}_{\ell+1} \tilde{s}_\ell = \tilde{s}_{\ell+1} \tilde{s}_\ell \tilde{s}_{\ell+1}$ is shown in [LP13, Theorem 6.6]. Indeed, these relations happen at a local part of the surface (in our case, torus) that looks like a cylinder.

On the other hand, the commutativity of the s_ℓ and the $\tilde{s}_\ell$ does not follow from [LP13, Theorem 12.2]. This is because in [LP13] we only considered the case when the pairs of parallel wires intersect once. In our case on the torus however, it is common to have horizontal and vertical wires intersect more than once: this happens for any $\mathfrak{d} \neq 1$. The proof in such a situation is essentially the same.

Indeed, if we have two pairs of parallel wires crossing as in Figure 10, but possibly more than once, we can realize each of the two corresponding R-moves by a sequence of local moves as above. It is a local check that performing one of the two sequences does not change the value of $p = \frac{\prod y_i - \prod x_i}{E(\boldsymbol{x}^{(j)}, \boldsymbol{y}^{(j)})}$ one needs to perform the other, because each of the $\prod x_i$, $\prod y_i$ and $E(\boldsymbol{x}^{(j)}, \boldsymbol{y}^{(j)})$ is not changed. It is also a local check that the two



Figure 10.

sequences commute once the parameters p for each are chosen, this is [LP13, Proposition 3.4]. Thus, commutativity follows.

5. Eigenvector map

In this section, we fix $f \in \mathcal{V}$ (see Definition 3.9) and consider the corresponding smooth curve C_f. For $j \in \mathbb{Z}$, we set $O_j := O_r$ if $j \equiv r \bmod N$. Similarly, for $j \in \mathbb{Z}$, we set $A_j := A_r$ if $j \equiv r \bmod m$.

5.1. **Generalities.** A divisor $D = \sum_i n_i P_i$ on an algebraic curve C is a finite formal integer linear combination of points P_i on C. We write $D \geq 0$ if $n_i \geq 0$, and say that D is *positive* in this case. The degree of D is given by $\deg(D) = \sum_i n_i$.

Given h a meromorphic function on C, we let $(h) = (h)_0 - (h)_\infty$ be the divisor of h. Here, $(h)_0$ denotes the divisor of zeroes, and $(h)_\infty$ denotes the divisor of poles. Two divisors D_1, D_2 are linearly equivalent, $D_1 \sim D_2$, if there exists a meromorphic function h such that $(h) = D_1 - D_2$. We write $[D]$ for the equivalence class of D with respect to linear equivalence. The Picard group $\mathrm{Pic}(C)$ is the abelian group of divisors on C modulo linear equivalence. For $j \in \mathbb{Z}$, we write $\mathrm{Pic}^j(C)$ for the part of the Picard group of C that has degree j, that is, $\mathrm{Pic}^j(C) := \{D : \text{ a divisor on } C \mid \deg(D) = j\}/\sim$.

To each divisor D we associate a space of meromorphic functions

$$\mathcal{L}(D) = \{f \mid (f) + D \geq 0\}.$$

If $D = \sum_i n_i P_i \geq 0$, then in words, $\mathcal{L}(D)$ consists of meromorphic functions which are allowed to have poles of order n_i at P_i. We have $\dim(\mathcal{L}(D_1)) = \dim(\mathcal{L}(D_2))$ if D_1 and D_2 are linearly equivalent.

5.2. **Positive divisors of a Lax matrix.** A divisor $D \in \mathrm{Pic}^g(C_f)$ of degree g is called *general* if $\dim \mathcal{L}(D) = 1$. A divisor $D \in \mathrm{Pic}(C_f)$ is called *regular* with respect to the points P and O_j if D is general and if $\dim \mathcal{L}(D + kP - \sum_{j=n-k}^{n} O_j) = 0$ for $k > 0$. For brevity, we will sometimes just say that a divisor is regular when it is regular with respect to P and the O_j.

Let us now fix $L(x) \in \beta^{-1}(f)$. Let

$$\Delta_{i,j} := (-1)^{i+j} |L(x) - y|_{i,j}$$

denote the (signed) (i,j)-th minor of the matrix $L(x)-y$.

Theorem 5.1 (cf. [vMM]). *There exists a positive divisor R of degree $2g+2$ supported on $S_{P,O} := \{P, O_1, \dots, O_N\}$, and uniquely defined positive general divisors $D_1, D_2, \dots, D_n, \bar{D}_1, \dots, \bar{D}_n$ of degree g such that for all $(i,j) \in [n]^2$, we have*

$$(\Delta_{i,j}) = D_j + \bar{D}_i + (j-i-1)P + \sum_{r=j+1}^{i-1} O_r - R.$$

In addition, $D_1, \dots, D_n$ have pairwise no common points, and $\bar{D}_1, \dots, \bar{D}_n$ have pairwise no common points.

See § 6.4 for the proof. Note that $\sum_{r=j}^{i} O_r$ is to be interpreted in a signed way:

$$\sum_{r=j}^{i} O_r := \sum_{r=j}^{\infty} O_r - \sum_{r=i+1}^{\infty} O_r = \begin{cases} \sum_{r=j}^{i} O_r & i \geq j, \\ 0 & i = j-1, \\ -\sum_{r=i+1}^{j-1} O_r & i \leq j-2. \end{cases}$$

For $L(x) \in \beta^{-1}(f)$, define $g_i := \Delta_{n,i}$. Since $L(x)-y$ is singular along C_f, the vector $g = (g_1, g_2, \dots, g_n)^\perp$ (thought of as a vector with entries that are rational functions on C_f) is an eigenvector of $L(x)-y$. We define g_i for $i \in \mathbb{Z}$ by $g_{i+n} = x^{-1} g_i$. We also define $h_i := g_i/g_n$ for $i \in \mathbb{Z}$. Thus $h_n = 1$ and $h_{i+n} = x^{-1} h_i$. The vector $(h_1, h_2, \dots, h_n)^\perp$ is also an eigenvector of $L(x)-y$.

Definition 5.2. For $L(x) \in \beta^{-1}(f)$, define the divisor $\mathcal{D} = \mathcal{D}(L(x))$ on C_f to be the minimum positive divisor satisfying

$$(h_i) + \mathcal{D} \geq \sum_{j=i+1}^{n} O_j - (n-i)P,$$

that is, $h_i \in \mathcal{L}(\mathcal{D} + (n-i)P - \sum_{j=i+1}^{n} O_j)$, for $i = 1, \dots, n-1$.

It follows from Theorem 5.1 that $\mathcal{D} = D_n$ from the theorem and that this divisor is uniquely determined by Definition 5.2. In particular, $\mathcal{D}$ is a positive regular divisor of degree g with respect to the points P and O_j, and we have

$$(h_i) = D_i - \mathcal{D} - (n-i)P + \sum_{j=i+1}^{n} O_j. \tag{5.1}$$

5.3. **The eigenvector map.** Let R_A (resp. R_O) denote the set of orderings of the m points $A_1, \dots, A_m$ (resp. $O_1, \dots, O_N$):

$$R_A := \{\nu(A_1, \dots, A_m) \mid \nu \in \mathfrak{S}_m\}, \qquad R_O := \{\tilde{\nu}(O_1, \dots, O_N) \mid \tilde{\nu} \in \mathfrak{S}_N\}.$$

By our assumption that $f \in \mathcal{V}$, the points A_r (resp. O_r) are distinct, so R_A has cardinality $m!$ and R_O has cardinality $N!$.

We write ν_r $(r = 1, \dots, m-1)$ (resp. $\tilde{\nu}_r$ $(r = 1, \dots, N-1)$) for the generator of $\mathfrak{S}_m$ (resp. $\mathfrak{S}_N$), which permutes the r-th and the $(r+1)$-st entry of $A \in R_A$ (resp. $O \in R_O$).

By our assumption that $f \in \mathcal{V}$, we have $f_c \neq 0$. Define $\mathcal{S}_f \subset (\mathbb{C}^*)^M$ by

$$\mathcal{S}_f = \left\{ (x_1, \dots, x_M) \in (\mathbb{C}^*)^M \mid \prod_{\ell=1}^{M} x_\ell = f_c \right\}.$$

For $\ell = 1, \ldots, M$, define c_ℓ to be the coefficient of the maximal power of x in the $(\ell, \ell+1)$ entry of $L(x)^{n/M}$. (Here, maximal power is the maximal power for a generic $L(x) \in \mathcal{L}$.) An explicit formula of c_ℓ for $L(x) \in \alpha(\mathcal{M})$ is given in Lemma 6.5.

We now define the *eigenvector map* η_f for $f \in \mathcal{V}$ by

$$\begin{aligned} \eta_f : \beta^{-1}(f) &\longrightarrow \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O && (5.2)\\ L(x) &\longmapsto ([\mathcal{D}], (c_1, \ldots, c_M), O), && (5.3)\end{aligned}$$

where the ordering $O = (O_1, \ldots, O_N)$ is uniquely determined from $(\Delta_{i,j})$ by Theorem 5.1. For $f \in \mathring{\mathcal{V}}$ we define the map ϕ_f by

$$\begin{aligned} \phi_f : \psi^{-1}(f) &\longrightarrow \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A && (5.4)\\ Q = (Q_1, \ldots, Q_m) &\longmapsto ([\mathcal{D}], (c_1, \ldots, c_M), O, A), && (5.5)\end{aligned}$$

where $A = (((-1)^n \epsilon_1, 0), \ldots, ((-1)^n \epsilon_m, 0))$, and $\epsilon_r(q)$ is given by (3.7). That $(c_1, \ldots, c_M)$ lies in $\mathcal{S}_f$ is the content of Lemma 6.6 below. We will omit the subscripts of η_f and ϕ_f and just write η and ϕ when no ambiguity will arise.

The following theorem should be compared to Theorem 1 in [vMM].

Theorem 5.3. *We have a one-to-one correspondence between $L(x) \in \beta^{-1}(\mathcal{V})$ and the following data:*

- (a) *$f \in \mathcal{V}$,*
- (b) *$([\mathcal{D}], c, O) \in \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O$ where $\mathcal{D}$ is a positive divisor on C_f of degree g, and regular with respect to P and the O_j.*

See §6.5 for the proof. It is shown in [LP11, Proof of Theorem 4.1] that the map $\alpha : \mathcal{M} \to \mathcal{L}$ is generically $m!$ to 1. Let $\mathring{\mathcal{M}}$ denote the open subset of $\mathcal{M}$ where (a) the map α is $m!$ to 1 (that is, $\mathring{\mathcal{M}} = \alpha^{-1}(\alpha(\mathring{\mathcal{M}})) \to \alpha(\mathring{\mathcal{M}})$ is a $m!$ to 1 map), and (b) the finite symmetric subgroup $\mathfrak{S}_m$ acting via the R-matrix is well-defined on $\mathring{\mathcal{M}}$.

Theorem 5.4. *Fix $f \in \mathring{\mathcal{V}}$. We have an injection from $\psi^{-1}(f) \cap \mathring{\mathcal{M}}$ to the collection of data $([\mathcal{D}], c, O, A) \in \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A$ such that*

- (a) *$\mathcal{D}$ is a positive divisor on C_f of degree g. It is regular with respect to P and O_j-s.*
- (b) *$c = (c_1, \ldots, c_M) \in \mathcal{S}_f$.*
- (c) *$O \in R_O$.*
- (d) *$A \in R_A$.*

See §6.6 for the proof. We will usually just write $\psi^{-1}(f)$ instead of $\psi^{-1}(f) \cap \mathring{\mathcal{M}}$ when no confusion arises (for example, when discussing the action of the R-matrix on the spectral data).

Let us perform a quick dimension count. The dimension of $\mathcal{M}$ or $\mathcal{L}$ is equal to mn. The dimension of $\mathrm{Pic}^g(C_f) \times \mathcal{S}_f$ is equal to $g + M - 1$. The number of lattice points in the convex full of $\{(k, 0), (0, n), (k+m, 0)\}$ is equal to $g + N + m + M$. Thus the dimension of $V_{n,m,k}$ (3.10) is equal to $g + N + m - 1$. Using (3.12) we obtain $\dim(\mathcal{M}) = \dim(\mathrm{Pic}^g(C_f) \times \mathcal{S}_f) + \dim(V_{n,m,k})$, consistent with Theorems 5.3 and 5.4.

6. Proofs from Section 5

The first three subsections are devoted to introduce key lemmas for Theorem 5.1 and 5.3.

6.1. Special zeroes and poles of the eigenvector.

Proposition 6.1. *The eigenvector* $g = (g_i)_i^T$ *of* $L(x) \in \beta^{-1}(f)$ *satisfies the following.*

(i) *For any* $j \in \mathbb{Z}$, *the rational function* g_j/g_{j+1} *on* C_f *has a zero of order one at* O_{j+1}.
(ii) *For any* $j \in \mathbb{Z}$, *the rational function* g_j/g_{j+1} *on* C_f *has a pole of order one at* P.

To prove this proposition, we shall need the following generalization of Theorem 3.2, whose proof is essentially the same as that of Theorem 3.2. Suppose $\mathbf{p} = \{p_1, p_2, \ldots, p_{n-1}\}$ is an unordered noncrossing family of n paths in G' using all the sources except i, and all the sinks except j. Identifying $[n] \setminus i \simeq [n-1] \simeq [n] \setminus j$ via the order-preserving bijection, we have that the non-abrupt paths in $\mathbf{p}$ induce a bijection of a subset $S \subset [n-1]$ with itself. We let $\mathrm{sign}(\mathbf{p})$ denote the sign of this permutation.

Let $|(L(x) - y)|_{i,j}$ denote the maximal minor of $(L(x) - y)$ complementary to the (i, j)-th entry.

Theorem 6.2. *We have*

$$|(L(x) - y)|_{i,j} = \sum_{\mathbf{p}=\{p_1,p_2,\ldots,p_{n-1}\}} \mathrm{sign}(\mathbf{p})\mathrm{wt}(p_1)\mathrm{wt}(p_2)\cdots\mathrm{wt}(p_{n-1}),$$

where the summation is over noncrossing (unordered) families of $n-1$ *paths in* G' *using all the sources except* i, *and all the sinks except* j. *In other words, the coefficient of* $x^a y^b$ *in* $|(L(x) - y)|_{i,j}$ *counts (with weights) families of highway paths that*

- *start at all sources but* i *and end at all sinks but* j;
- *do not cross each other;*
- *cross the* x*-line exactly* a *times;*
- *contain exactly* b *abrupt paths and* $n - b - 1$ *non-abrupt paths.*

Example 6.3. Let $(n, m, k) = (3, 3, 1)$. The Lax matrix is given as follows: $L(x) =$

$$\begin{pmatrix} q_{1,1}x + q_{2,3}x + q_{3,2}x & q_{1,1}q_{2,1}x + q_{1,1}q_{3,3}x + q_{2,3}q_{3,3}x & q_{1,1}q_{2,1}q_{3,1}x + x^2 \\ q_{1,2}q_{2,2}q_{3,2} + x & q_{1,2}x + q_{2,1}x + q_{3,3}x & q_{1,2}q_{2,2}x + q_{1,2}q_{3,1}x + q_{2,1}q_{3,1}x \\ q_{1,3}q_{2,3} + q_{1,3}q_{3,2} + q_{2,2}q_{3,2} & q_{1,3}q_{2,3}q_{3,3} + x & q_{1,3}x + q_{2,2}x + q_{3,1}x \end{pmatrix}$$

Consider the minor $|(L(x) - y)|_{1,2} =$ with rows $2, 3$ and columns $1, 3$. Here are some terms that appear in it:

$$|(L(x) - y)|_{1,2} = q_{2,2}x^2 - q_{1,3}q_{2,3}q_{1,2}q_{2,2}x - xy - \cdots .$$

The term $q_{2,2}x^2$ for example is formed by two paths: one starting at source 3, turning at $q_{1,3}$, turning at $q_{1,2}$, going through at $q_{2,2}$, turning at $q_{3,2}$, turning at $q_{3,1}$ and thus finishing in sink 3; the other a staircase path starting at source 2, turning at each of $q_{1,2}$, $q_{1,1}$, $q_{2,1}$, $q_{2,3}$, $q_{3,3}$, $q_{3,2}$ and finishing at sink 1. The first path contributes weight $q_{2,2}x$, while the second contributes weight x. Note that source 1 and sink 2 remain unused, as they should according to the theorem. Also note the the paths give bijection $2 \mapsto 1$

and $3 \mapsto 3$ between the used sources and sinks. The induced permutation is the identity permutation, and that is why we have sign $+$ in front of this term.

The term $-xy$ corresponds to one abrupt path of weight $-y$ from source 3 to sink 3, and one staircase path, the same as for the previous term. The term $-q_{1,3}q_{2,3}q_{1,2}q_{2,2}x$ corresponds to two paths that induce the map $2 \mapsto 3$ and $3 \mapsto 1$ on sources and sinks, of weights $q_{1,2}q_{2,2}x$ and $q_{1,3}q_{2,3}$ respectively. The minus sign arises since the induced permutation in S_2 is a transposition.

Let us prove Proposition 6.1. (i) If $N = 1$, the result is easy: g_j/g_{j+n} vanishes to order n at O_1. We also have a shift automorphism $s : \mathcal{L} \to \mathcal{L}$ (see the proof of Proposition 6.1(ii)) which sends O_1 to O_1 and pulls g_j/g_{j+1} back to g_{j+1}/g_{j+2}. We will thus assume $N > 1$.

For each i, define

$$v^{(i)} := \left((-1)^1|L(x) - y|_{i,1}, \ldots, (-1)^n|L(x) - y|_{i,n}\right)^T.$$

We shall think of $v^{(i)}$ as a vector whose entries lie in the coordinate ring of the affine plane curve $\tilde{C}_f$. Like the vector $g = (g_1, g_2, \ldots, g_n)^T$, the vectors $v^{(i)}$ are (nonzero) eigenvectors of the matrix $L(x)$. The matrix $L(x)$ is singular along the curve $\tilde{C}_f$, but generically it has rank $n - 1$. Thus for any i, the vectors g and $v^{(i)}$ are multiples of each other. More precisely, $g/v^{(i)}$ is a rational function on C_f. To show that g_j/g_{j+1} has a zero at O_{j+1}, we shall calculate using a convenient choice of $v^{(i)}$.

Choose $v = v^{(j)}$. Then $v_j = |L(x) - y|_{j,j}$ and $v_{j+1} = |L(x) - y|_{j,j+1}$. Thus, v_j counts the families of paths that start at all sources but j and end at all sinks but j, as in Theorem 6.2.

Let us make the substitution $(x, y) \sim (u^{n'}, -(-\sigma_j)^{1/n'}u^{k'})$ inside v_j, and let $v'_j(x, y)$ denote the terms of in $v_j(x, y)$ that give the lowest degree in u after the substitution. Call this degree d. The terms in $v_j(x, y)$ are obtained by either taking abrupt paths, or by taking coils. In the proof of Lemma 3.5 we defined the N coils in the network G, which we denote $C_1, C_2, \ldots, C_N$. To obtain a path family $\mathbf{p}$ contributing to v'_j, instead of C_j, we include the $n' - 1$ abrupt paths which use the vertices j' where $j \neq j'$ but $j \equiv j'$ mod N. For each of the coils C_t where $t \neq j$, we can either include the coil (that is, include the n' paths in G') in $\mathbf{p}$, or we use the corresponding n' abrupt paths instead. In particular, considering C_{j+1} we see that v'_j has a factor of $(\sigma_{j+1}x^{k'} + (-y)^{n'})$. This factor vanishes under the substitution $(x, y) \sim (u^{n'}, -(-\sigma_{j+1})^{1/n'}u^{k'})$, thus creating a zero at the point O_{j+1} of order at least $d + 1$.

For v_{j+1}, we count families of paths that start at all sources but j and end at all sinks but $j + 1$. Let $v'_{j+1}(x, y)$ denote the terms of lowest degree in $v_{j+1}(x, y)$. This lowest degree is again equal to d as well (we caution that if $j = n$, then we take $v_{j+1} := x^{-1}v_1$). The calculation of $v'_{j+1}(x, y)$ is similar to that of $v_j(x, y)$, except that instead of a single "incomplete" coil with index j modulo N, we have two incomplete coils C_j and C_{j+1} with indices j and $j + 1$ modulo N. We obtain

$$v'_{j+1}(x, y) = a(q)x^{\alpha}y^{\beta} \prod_{t \neq j, j+1} (\sigma_t x^{k'} + (-y)^{n'})$$

where $a(q)$ is some nonzero polynomial in q_{ab}-s, and α, β are nonnegative integers, and the product is over $t \in \{1, 2, \dots N\}$ not equal to j or $j+1$ modulo N. As a result, we see that v_{j+1} vanishes at O_{j+1} of order exactly d.

Thus g_j/g_{j+1} has a zero at O_{j+1}. The fact that g_j/g_{j+1} vanishes to exactly order 1 follows since we know that g_j/g_{j+n} vanishes to the order of n' at each O_i.

(ii) For $L(x) \in \beta^{-1}(f)$, we have rational functions g_i/g_{i+1} on C_f. To emphasize the dependence of g_i/g_{i+1} on L, we write $g_i/g_{i+1}(L)$. Let $\varsigma^* : \mathcal{L} \to \mathcal{L}$ be the shift map given by $L(x) \mapsto P(x)L(x)P(x)^{-1}$. (We will introduce the shift operator ς on $\mathcal{M}$ in § 7.2.) Then $\beta(\varsigma^*(L(x))) = \beta(L(x))$ is invariant under r. Let Y be a Zariski dense subset of $\beta^{-1}(\mathring{\mathcal{V}})$ such that $\varsigma^*(Y) = Y$. Then there exists a nonnegative integer a_i given by

$$a_i = \max\{\mathrm{ord}_P(g_i/g_{i+1}(L))_\infty \mid L \in Y\},$$

where $\mathrm{ord}_P(g_i/g_{i+1}(L))_\infty$ denotes the order of the pole of $g_i/g_{i+1}(L)$ at P. Since $g_i/g_{i+1}(\varsigma^*(L)) = g_{i-1}/g_i(L)$ as rational function on $C_{\beta(L)}$, we conclude that for all i, we have $\mathrm{ord}_P(g_i/g_{i+1}(L))_\infty = a := \min_i\{a_i\}$. But we know that $\mathrm{ord}_P(g_i/g_{i+n}(L))_\infty = n$ for all i and $L \in \mathcal{L}$. Thus we must have $a = 1$. Since this value does not depend on the choice of Y, the claim follows.

6.2. **A holomorphic differential.** Let ζ be the differential form on the curve C_f given by

$$\zeta = \frac{x^{k-1}dy}{\frac{\partial f}{\partial x}}.$$

Lemma 6.4. *The divisor of the differential form ζ is supported on $S_{P,O} = \{P, O_1, \dots, O_N\}$:*

$$(\zeta) = (k'-1)\sum_{i=1}^{N} O_i + ((n-1)m - k - M)\, P.$$

When $g \geq 1$, it is holomorphic.

Proof. By using local expansions of f around O_i and P, we get

$$\left(\frac{\partial f}{\partial x}\right) = (n' - k'n)\sum_{i=1}^{N} O_i + n(k+m-1)P,$$

$$(dy) = (k'-1)\sum_{i=1}^{N} O_i - (m+k+M)P.$$

From these and (x) in §4.3, we obtain (ζ).

Now we prove that if $g \geq 1$ then $(n-1)m - k - M \geq 0$. Let p be the number of integer points inside a parallelogram in $\mathbb{R}^2$, whose vertices are $(0,n)$, $(k,0)$, $(k+m,0)$ and (m,n). We have $p = 2g + M - 1$, since the parallelogram is composed of the Newton polygon $N(f)$ and its copy, sharing the upper hull of $N(f)$. Then the claim is equivalent to that if $g \geq 1$, then $p + N - 1 \geq M + k$.

When $k = n$, $N = n$ follows. Then we have $p + N - 1 - (M+k) = p - (M+1)$. It reduces to $2g - 2$, which is non-negative when $g \geq 1$.

When $m + k = n$, $M = n$ follows. Then we have $g = \frac{1}{2}(n(m-1) - m - N + 1)$ which is negative if $m = 1$. So we assume $m \geq 2$. We have $p + N - 1 = m(n-1)$ and $M + k = 2n - m$. Thus, we obtain $p + N - 1 - (M+k) \geq 0$ when $m \geq 2$.

When $n > k$, we prove the claim by choosing $M + k$ different points from p points in the parallelogram. When $m + k < n$ (resp. $m + k > n$), we can choose $M - 1$ points on the upper hull of $N(f)$, k points; $(i, n - i)$ for $i = 1, \dots, k$ inside the upper (resp. lower) triangle, and one point inside the lower (resp. upper) triangle.

Finally the claim follows. □

6.3. **About** $(c_1, \dots, c_M) \in \mathcal{S}_f$**.** Recall that c_ℓ is the coefficient of the maximal power of x in the $(\ell, \ell + 1)$ entry of $L(x)^{n/M}$ defined in §5.3.

Lemma 6.5. *When* $L(x) = \alpha(q = (q_{ij}))$*, we have*

$$c_\ell = \sum_{i+j\equiv \ell+1 \mod M} q_{ij}.$$

Proof. By Lemma 4.1, it suffices to prove the lemma for $f(x, y) \in \psi(\mathcal{M})$. We apply Lemma 3.1 to interpret the c_ℓ as counting almost snake paths in G, that is, paths that turn at all steps but one. The almost snake paths p that are counted start at vertex ℓ and end at vertex $\ell + 1$, and when drawn in the cylindric network G', they are disjoint unions of n/M paths. In other words, p wraps around the picture Figure 4 horizontally n/M times. Such paths p are completely determined by the single vertex that the path goes through. With ℓ fixed, the weights of these vertices are exactly the $q_{i,j}$ with $i + j \equiv \ell + 1 \mod M$. □

Recall that we defined f_c in (3.5).

Lemma 6.6. *When* $L(x) \in \beta^{-1}(f)$*, we have*

$$\prod_{\ell=1}^{M} c_\ell = f_c = \pm \sum_{(j,i)\in L_c} f_{i,j}. \tag{6.1}$$

Thus the product of all c_ℓ *is constant on* $\psi^{-1}(f)$*.*

Proof. We first show that

$$\text{degree of } f_{i,j} \text{ in the } q_{r,s} = (m + k)n - ((m + k)i + nj). \tag{6.2}$$

Indeed, lift a family $\mathbf{p}$ of paths that contributes to $f_{i,j}$ (as in Theorem 3.2) to the universal cover, as in Figure 6. By Theorem 3.2, the family $\mathbf{p}$ includes exactly i abrupt paths and $n - i$ non-abrupt paths. Let us suppose that the sources used by the $n - i$ non-abrupt paths are $(1/2, z_1), \dots, (1/2, z_{n-i})$ and the sinks are $(m + 1/2, w_1), \dots, (m + 1/2, w_{n-i})$, where $1 \le z_r \le n$ and $1 \le w_r \le n + m$. To agree with our convention in Figure 4 that source and sink labels increase as we go down, we will take the y-coordinate to increase as we go down in Figure 6; but otherwise, the coordinates we are using are the usual Cartesian coordinates.

For each r, let $w'_r \in [1 - k, n - k]$ be chosen so that $w'_r \equiv w_r \mod n$. From Theorem 3.2 it also follows that

$$\sum_{r=1}^{n-i} (w_r - w'_r)/n = i,$$

as this is the number of times the x-line would be crossed by the paths. Also, we know that $\{w_1, \dots, w_{n-i}\} = \{z_1 - k, \dots, z_{n-i} - k\} \mod n$ and thus $\{w'_1, \dots, w'_{n-i}\} =$

$\{z_1 - k, \ldots, z_{n-i} - k\}$. This is because our collection of non-abrupt paths must form a collection of cycles in G. The number of the $q_{r,s}$ a path from $(1/2, z_r)$ to $(m + 1/2, w_s)$ picks up is equal to $z_r + m - w_s$, and summing over all the non-abrupt paths in $\mathbf{p}$ we get

$$\begin{aligned}\text{degree of } f_{i,j} \text{ in the } q_{r,s} &= m(n-j) + \sum_{r=1}^{n-i} z_r - \sum_{s=1}^{n-i} w_s \\ &= m(n-j) + \sum_{r=1}^{n-i} z_r - \sum_{s=1}^{n-i} w'_s + jn \\ &= m(n-i) + (n-i)k - jn = (m+k)n - ((m+k)i + nj),\end{aligned}$$

as claimed. This degree is equal to M when $(j, i) \in L_c$.

Now we prove the lemma. Let us partition all the edges in the network G into M snake paths, $H_1, \ldots, H_M$. Suppose p is a closed path in G. Then we can partition p into *snake intervals* – maximal contiguous segments of p that follow one of the H_ℓ. Such snake intervals are connected by a *horizontal step* going through a vertex (and picking up its weight), and moving from a snake path H_ℓ to the next snake path $H_{\ell+1}$. It is easy to see that the vertices which separate H_ℓ from $H_{\ell+1}$ are exactly the ones labelled with q_{ij} where $i + j \equiv \ell + 1 \mod M$.

In order for p to be closed, it must consist of c horizontal steps, where c is a multiple of M, since p must come back to the snake path it started at. When $(j, i) \in L_c$, (6.2) shows that all families of toric paths that contribute to $f_{i,j}$ consist of just a single closed path in G that picks up exactly one of the q_{ij} from each of the sets $T_\ell = \{q_{ij} \mid i + j \equiv \ell + 1 \mod M\}$, for $\ell = 1, \ldots, M$. Using Lemma 6.5, we see that each such term appears exactly once in the product $\prod_{\ell=1}^{M} c_\ell$.

We need to show that this is not only an injection, but a bijection. For each term contributing to $\prod_{\ell=1}^{M} c_\ell$, it suffices to find a closed path in G that has this weight. By (6.2), such a closed path would necessarily contribute to $f_{i,j}$ for $(j, i) \in L_c$.

For each $q_{r,s}$ appearing in our chosen term of $\prod_{\ell=1}^{M} c_\ell$, we draw a horizontal step through the vertex labelled $q_{r,s}$ in the network G. From the endpoint of each such horizontal step (say from H_ℓ to $H_{\ell+1}$), attach a snake interval in $H_{\ell+1}$ leading to the start point of the horizontal step which goes from $H_{\ell+1}$ to $H_{\ell+2}$. Some of these snake intervals may be empty. When we glue everything together, we get the desired closed path. □

6.4. **Sketch proof of Theorem 5.1.** The proof is analogous that of [vMM]. We explain the differences. When $n - k \le m \le 2n - k$, the triple of integers (N, M, M') in van Moerbeke and Mumford's work [vMM] corresponds to the parameters $(n, n-k, m+k-n)$ in our case.

One first shows that there is a positive regular divisor of degree g, denoted $\mathcal{D}$, satisfying Definition 5.2. The correspondence $L(x) \mapsto \mathcal{D}$ is the main construction in [vMM, Theorem 1]. Their results do not formally apply since our Lax matrix is not *regular* in their terminology. Nevertheless, the properties of $L(x) \mapsto \mathcal{D}$ is proved in the same manner as [vMM, Lemmas 3 and 4], where Proposition 6.1 takes the place of [vMM, Lemma 2].

The divisors D_i and $\bar{D}_j$ of Theorem 5.1 are obtained by shifting and transposing the Lax matrix. [vMM, Proposition 1] then says, in our terminology,

$$(\Delta_{i,j}\zeta) = D_j + D'_i + (j-i-1)P + \sum_{r=j+1}^{i-1} O_r$$

where ζ is the differential of Lemma 6.4. We now take $R = (\zeta)$ to obtain Theorem 5.1. The statement about common points is on [vMM, p.117].

6.5. **Proof of Theorem 5.3.** We shall also use the following lemma which is proved in a similar way to [vMM, Lemma 5].

Lemma 6.7. *Suppose $\mathcal{D}$ is a positive divisor on C_f of degree g. which is regular with respect to the points P and O_j. We have*

(i) $\dim \mathcal{L}(\mathcal{D} + (n-i)P - \sum_{j=i+1}^{n} O_j) = 1$ *for* $i \in \mathbb{Z}$.

(ii) *Suppose for each $i \in \mathbb{Z}$, we have fixed a nonzero element $h_i \in \mathcal{L}(\mathcal{D} + (n-i)P - \sum_{j=i+1}^{n} O_j)$. Then for $i_1 \le i_2$, and any $h \in \mathcal{L}(\mathcal{D} + (n-i_1)P - \sum_{j=i_2+1}^{n} O_j)$, there are unique scalars $b_{i_1}, \dots, b_{i_2}$ such that*

$$h = \sum_{i=i_1}^{i_2} b_i h_i,$$

with $b_{i_1}, b_{i_2} \neq 0$.

Let us prove the theorem. Due to Theorem 5.1, Lemma 6.6 and the assumption on $\mathcal{V}$, for $L(x) \in \beta^{-1}(\mathcal{V})$ we have $f := \beta(L(x)) \in \mathcal{V}$, and $\eta_f(L(x)) = ([\mathcal{D}], c, O)$ satisfying (b). In the following we construct the inverse of η_f (5.2) for $f \in \mathcal{V}$. Around $P \in C_f$ we have the local expansion $(x, y) = (x_0 u^{-n}, y_0 u^{-m-k})$ as $u \to 0$. We take $c = (c_1, \dots, c_M) \in \mathcal{S}_f$, and recursively define d_i ($i \in \mathbb{Z}$) by

$$d_n = 1, \qquad d_i = c_\ell \, d_{i+1} \quad \text{for } i \equiv \ell \mod M.$$

Let $h_i \in \mathcal{L}(\mathcal{D} + (n-i)P - \sum_{j=i+1}^{n} O_j)$ have an expansion around P as $h_i = d_i u^{-n+i} + \cdots$.

Since $y\, h_i \in \mathcal{L}(\mathcal{D} + (n+m+k-i)P - \sum_{j=-k+i+1}^{n} O_j)$, using Lemma 6.7, we have unique scalars $b_{i,j}$ satisfying

$$y\, h_i = \sum_{j=n-m-k}^{n-k} x\, b_{i,i+j} h_{i+j}, \qquad b_{i,i-m-k}, b_{i,i-k} \neq 0.$$

By expanding it around P, we get $b_{i,i-m-k} = y_0 d_i / d_{i-m-k} = y_0 / f_c^{\frac{m+k}{M}}$ independent of i. Define an infinite matrix $A = (a_{ij})_{i \in [n], j \in \mathbb{Z}}$ by

$$a_{i,i+j} = \begin{cases} b_{i,i+j} \dfrac{f_c^{\frac{m+k}{M}}}{y_0} & \text{for } -m-k \le j \le -k, \\ 0 & \text{otherwise.} \end{cases}$$

By using (3.3) for this A, we obtain $L(A; x) \in \beta^{-1}(f)$.

6.6. **Proof of Theorem 5.4.** Fix $L(x) \in \beta^{-1}(f)$, and assume that $\alpha^{-1}(L)$ contains $Q \in \mathring{\mathcal{M}}$. By Lemma 7.1, the action of the finite symmetric group $\mathfrak{S}_m$ via the R-matrix action of §2.2 preserves the Lax matrix $L(x)$. (See §7 for more discussion of this action.) Thus $\mathfrak{S}_m \cdot Q \subseteq \alpha^{-1}(L)$. By (2.1), the $\mathfrak{S}_m$ action induces the permutation action on the m quantities $\epsilon_1(q), \ldots, \epsilon_m(q)$, which corresponds exactly to the permutation action on R_A. Also, for $f \in \mathring{\mathcal{V}}$, these m quantities $\epsilon_1(q), \ldots, \epsilon_m(q)$ are distinct. So $|\mathfrak{S}_m \cdot Q| = m! = |\alpha^{-1}(L)|!$ implying that $\mathfrak{S}_m \cdot Q = \alpha^{-1}(L)$. Thus the map $\alpha^{-1}(L) \to (L, R_A)$ is an injection, and the claim follows.

7. Actions on $\mathcal{M}$

We study the family of commuting extended affine symmetric group actions on $\mathcal{M}$, introduced in § 2.2. We also study another family of actions on $\mathcal{M}$, which we call the *snake path actions*. The main result in this section is Theorem 7.3.

Throughout §7–9, we fix $f \in \mathring{\mathcal{V}}$. For simplicity, we write $Q_i := Q_i(x)$ in the rest of this section.

7.1. The behaviour of spectral data under the extended affine symmetric group actions.

Recall the definitions of energy and R-matrix from §2.1. Let $A(x)$ and $B(x)$ be two n by n matrices of the shape (3.1), with diagonal entries $\mathbf{a} = (a_1, \ldots, a_n)$ and $\mathbf{b} = (b_1, \ldots, b_n)$. Define the *energy* by $E(A, B) := E(\mathbf{a}, \mathbf{b})$ and the *R-matrix* to be the transformation $(A(x), B(x)) \mapsto (B'(x), A'(x))$, where $B'(x)$ and $A'(x)$ have diagonal entries $\mathbf{b}'$ and $\mathbf{a}'$ respectively, and $R(\mathbf{a}, \mathbf{b}) = (\mathbf{b}', \mathbf{a}')$. The following is proved for example in [LP13, Corollary 6.4].

Lemma 7.1. *Suppose $R(A(x), B(x)) = (B'(x), A'(x))$. Then we have $A(x)B(x) = B'(x)A'(x)$.*

Indeed, Lemma 7.1 and the condition that R is non-trivial uniquely determine the R-matrix as a birational transformation.

Now we reformulate the $W \times \tilde{W}$ actions introduced in § 2.2, in our current terminology. The action of the extended affine symmetric group W is generated by s_i $(1 \leq i \leq m-1)$ and π, where for $1 \leq i \leq m-1$, s_i acts by

$$s_i : (Q_1, Q_2, \ldots, Q_m) \longmapsto (Q_1, \ldots, Q'_{i+1}, Q'_i, \ldots, Q_m)$$

where (Q'_{i+1}, Q'_i) is the R-matrix image of (Q_i, Q_{i+1}). The operator π acts as

$$\pi : (Q_1, Q_2, \ldots, Q_m) \longmapsto (Q_2, Q_3, \ldots, Q_m, P(x)^k Q_1 P(x)^{-k}).$$

The operator e_u acts by moving, via the R-matrix, the last u terms of

$$(Q_{u+1}, \ldots, Q_m, P(x)^k Q_1 P(x)^{-k}, \ldots, P(x)^k Q_u P(x)^{-k})$$

to the first u positions, keeping them in order.

Recall from §2.2 that we also have $\tilde{Q} = (\tilde{Q}_1, \dots, \tilde{Q}_N) = (\tilde{q}_{i,j})$ coordinates on $\mathcal{M}$, where $\tilde{Q}_i := \tilde{Q}_i(x)$ is an mn' by mn' matrix:

$$\tilde{Q}_i(x) = \begin{pmatrix} \tilde{q}_{i,1} & 0 & 0 & x \\ 1 & \tilde{q}_{i,2} & 0 & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & 0 & 1 & \tilde{q}_{i,mn'} \end{pmatrix}.$$

Similarly, the action of $\tilde{W}$ generated by $\tilde{s}_r$ $(1 \le r \le N-1)$ and $\tilde{\pi}$ on $\tilde{Q} = (\tilde{Q}_1, \dots, \tilde{Q}_N) \in \mathcal{M}$ is described as follows: for $1 \le r \le N-1$, $\tilde{s}_r$ acts by

$$\tilde{s}_r : (\tilde{Q}_1, \dots, \tilde{Q}_N) \longmapsto (\tilde{Q}_1, \dots, \tilde{Q}'_{r+1}, \tilde{Q}'_r, \dots, \tilde{Q}_N)$$

where $(\tilde{Q}'_r, \tilde{Q}'_{r+1})$ is the R-matrix image of $(\tilde{Q}_{r+1}, \tilde{Q}_r)$. The operator $\tilde{\pi}$ acts as

$$\tilde{\pi} : (\tilde{Q}_1, \tilde{Q}_2, \dots, \tilde{Q}_N) \longmapsto (\tilde{Q}_2, \tilde{Q}_3, \dots, \tilde{Q}_N, \tilde{P}(x)^{m\bar{k}'} \tilde{Q}_1 \tilde{P}(x)^{-m\bar{k}'}),$$

where $\tilde{P}(x)$ is the mn' by mn' version of $P(x)$ (3.4).

Recall that we denote by $\mathfrak{S}_m \subset W$ (resp. $\mathfrak{S}_N \subset \tilde{W}$) the finite symmetric group generated by $s_1, \dots, s_{m-1}$ (resp. $\tilde{s}_1, \dots, \tilde{s}_{N-1}$). Then W is generated by $\mathfrak{S}_m$ and π, and $\tilde{W}$ is generated by $\mathfrak{S}_N$ and $\tilde{\pi}$.

Recall that in § 2.4 we define divisors $\mathcal{O}_u$ and $\mathcal{A}_u$ on C_f as follows:

$$\mathcal{O}_u = uP - \sum_{i=N-u+1}^{N} O_i, \qquad \mathcal{A}_u = uP - \sum_{i=1}^{u} A_i, \tag{7.1}$$

for $u \in \mathbb{Z}_{\ge 0}$. Let τ acts on $\mathcal{S}_f$ by $(c_1, \dots, c_M) \mapsto (c_M, c_1, \dots, c_{M-1})$.

Theorem 7.2. (i) *The actions of* $\mathfrak{S}_m$ *and* π *on* $\psi^{-1}(f)$ *induce the following transformations on* $\mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A$*: for* $r = 1, \dots, m-1$, s_r *induces*

$$([\mathcal{D}], c, O, A) \mapsto ([\mathcal{D}], c, O, \nu_r(A)), \tag{7.2}$$

and π *induces*

$$([\mathcal{D}], c, O, A) \mapsto ([\mathcal{D} - \mathcal{A}_1], \tau^{-1}(c), O, \nu_{m-1}\nu_{m-2} \cdots \nu_1(A)). \tag{7.3}$$

(ii) *The actions of* $\mathfrak{S}_N$ *and* $\tilde{\pi}$ *on* $\psi^{-1}(f)$ *induce the following transformations on* $\mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A$*: for* $r = 1, \dots, N-1$, $\tilde{s}_r$ *induces*

$$([\mathcal{D}], c, O, A) \mapsto ([\mathcal{D}], c, \tilde{\nu}_{N-r}(O), A), \tag{7.4}$$

and $\tilde{\pi}$ *induces*

$$([\mathcal{D}], c, O, A) \mapsto ([\mathcal{D} + \mathcal{O}_1], \tau(c), \tilde{\nu}_1 \cdots \tilde{\nu}_{N-1}(O), A). \tag{7.5}$$

Theorems 7.2 will be proved in §7.5. The following theorem states that the commuting $\mathbb{Z}^m$ and $\mathbb{Z}^N$ actions on $\mathrm{Pic}^g(C_f)$ are linearized by the spectral map ϕ.

Theorem 7.3. *The following diagrams are commutative:*

$$\begin{tikzcd} \psi^{-1}(f) \arrow[r, "\phi"] \arrow[d, "e_u"'] & \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A \arrow[d, "{(-[\mathcal{A}_u], \tau^{-u}, id, id)}"] \\ \psi^{-1}(f) \arrow[r, "\phi"'] & \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A, \end{tikzcd} \tag{7.6}$$

for $u = 1, \dots, m$, *and*

$$\begin{tikzcd}
\psi^{-1}(f) \arrow[r, "\phi"] \arrow[d, "\tilde{e}_u"'] & \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A \arrow[d, "{(+[\mathcal{O}_u], \tau^u, id, id)}"] \\
\psi^{-1}(f) \arrow[r, "\phi"'] & \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A,
\end{tikzcd} \tag{7.7}$$

for $u = 1, \dots, N$. *Here* $-[\mathcal{A}_u]$ *and* $+[\mathcal{O}_u]$ *respectively act on* $\mathrm{Pic}^g(C_f)$ *as* $[\mathcal{D}] \mapsto [\mathcal{D} - \mathcal{A}_u]$ *and* $[\mathcal{D}] \mapsto [\mathcal{D} + \mathcal{O}_u]$.

Proof. We shall show (7.6). The proof of (7.7) is done in the similar way by exchanging the rules of the special points $\{A_i\}_{i \in [m]}$ for that of $\{O_j\}_{j \in [N]}$.

From (7.2) and (7.3), it follows that a non-trivial part is to show e_u (2.8) induces the action on $\mathrm{Pic}^g(C_f) \times R_A$:

$$([\mathcal{D}], A) \mapsto ([\mathcal{D} - \mathcal{A}_u], A).$$

Note that for $1 \le i < j \le m$, $\nu_{j-1}\nu_{j-2} \cdots \nu_i \in \mathfrak{S}_m$ acts on R_A as

$$(A_1, \dots, A_m) \mapsto (A_1, \dots, A_{i-1}, A_{i+1}, \dots, A_j, A_i, A_{j+1}, \dots, A_m). \tag{7.8}$$

Thus, (7.3) indicates that π induces $([\mathcal{D}], (A_1, \dots, A_m)) \mapsto ([\mathcal{D} - \mathcal{A}_1], (A_2, \dots, A_m, A_1))$, and π^u induces

$$([\mathcal{D}], (A_1, \dots, A_m)) \mapsto ([\mathcal{D} - \mathcal{A}_u], (A_{u+1}, \dots, A_m, A_1, A_2, \dots, A_m)).$$

Further, (7.2) denotes that s_r does not change a point in $\mathrm{Pic}^g(C_f)$ and acts on R_A as a permutation $\nu_r \in \mathfrak{S}_m$. The rest part $(s_u \cdots s_{m-1})(s_{u-1} \cdots s_{m-2}) \cdots (s_1 \cdots s_{m-u})$ of e_u changes $(A_{u+1}, \dots, A_m, A_1, A_2, \dots, A_m) \in R_A$ following (the inverse of) (7.8) as

$$\begin{aligned}
&(A_{u+1}, \dots, A_m, A_1, A_2, \dots, A_m) \overset{\nu_1 \cdots \nu_{m-u}}{\mapsto} (A_1, A_{u+1}, \dots, A_m, A_2, \dots, A_m) \\
&\overset{\nu_2 \cdots \nu_{m-u+1}}{\mapsto} \cdots \overset{\nu_{u-1} \cdots \nu_{m-2}}{\mapsto} (A_1, \dots, A_{u-1}, A_{u+1}, \dots, A_m, A_u) \overset{\nu_u \cdots \nu_{m-1}}{\mapsto} (A_1, \dots, A_m). \qquad \square
\end{aligned}$$

We remark that e_u and $\tilde{e}_u$ act on the sets R_A and R_O as the identity transformations.

7.2. **The shift operator.**

We define a shift operator ς acting on $\mathcal{M}$ as

$$\varsigma : (Q_i)_{1 \le i \le m} \mapsto (P(x) Q_i P(x)^{-1})_{1 \le i \le m}. \tag{7.9}$$

It is easy to see that it induces an action $\varsigma^* : \mathcal{L} \to \mathcal{L}$ given by $\varsigma^* : L(x) \mapsto P L(x) P^{-1}$. The following result generalizes [I08, Proposition 2.6].

Proposition 7.4. *The following diagram is commutative:*

$$\begin{tikzcd}
\psi^{-1}(f) \arrow[r, "\phi"] \arrow[d, "\varsigma"'] & \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A \arrow[d, "{(+[P - O_N], \tau, \tilde{\nu}_1 \cdots \tilde{\nu}_{N-1}, id)}"] \\
\psi^{-1}(f) \arrow[r, "\phi"'] & \mathrm{Pic}^g(C_f) \times \mathcal{S}_f \times R_O \times R_A,
\end{tikzcd} \tag{7.10}$$

where $+[P - O_N]$ *acts on* $\mathrm{Pic}^g(C_f)$ *as* $[\mathcal{D}] \mapsto [\mathcal{D} + P - O_N]$.

Proof. Suppose $Q \in \psi^{-1}(f)$ and $\phi(Q) = ([\mathcal{D}], c, O, A)$. Then $\phi \circ \varsigma(Q) = ([\mathcal{D}'], \tau(c), \tilde{\nu}_1 \cdots \tilde{\nu}_{N-1}(O), A)$, where $\mathcal{D}'$ is some positive divisor of degree g. We set $(O'_1, \dots, O'_N) := \tilde{\nu}_1 \cdots \tilde{\nu}_{N-1}(O_1, \dots, O_N) = (O_N, O_1, \dots, O_{N-1})$.

Recall that $g = (g_1, g_2, \ldots, g_n)^T$ denotes the eigenvector of $\alpha(Q)$. An eigenvector of $\alpha \circ \varsigma(Q)$ is $(g'_1, g'_2, \ldots, g'_n)^T := (xg_n, g_1, g_2, \ldots, g_{n-1})^T$. Define $h'_i := g'_i/g'_n = g_{i-1}/g_{n-1} = h_{i-1}/h_{n-1}$; these ratios do not depend on which eigenvector of $\alpha \circ \varsigma(q)$ we chose. Then by (5.1)

$$\begin{aligned}(h'_i) = (h_{i-1}) - (h_{n-1}) &= (D_{i-1} - \mathcal{D} - (n-i+1)P + \sum_{j=i}^{n} O_j) - (D_{n-1} - \mathcal{D} - P + O_n) \\ &= D_{i-1} - D_{n-1} - (n-i)P + \sum_{j=i+1}^{n} O'_j.\end{aligned}$$

By the uniqueness of $\mathcal{D}'$ it follows that the divisor $\mathcal{D}'$ (resp. D'_i) is equal to D_{n-1} (resp. D_{i-1}). Furthermore, $\mathcal{D}' - \mathcal{D} \sim P - O_n$, as required.

□

7.3. $W \times \tilde{W}$ **actions on Lax-matrix.** For $r = 1, \ldots, N-1$, $i = 0, \ldots, n'-1$ and $s = 0, \ldots, m-1$, using the energy in § 2.1 define

$$E^{(s)}_{r+ki} := E(\tilde{P}(x)^{-mi-s}\tilde{Q}_r\tilde{P}(x)^{mi+s}, \tilde{P}(x)^{-mi-s}\tilde{Q}_{r+1}\tilde{P}(x)^{mi+s}).$$

Define an n by n matrix $B_{s,r}$ by:

$$B_{s,r} = \mathbb{I}_n + \sum_{i=0}^{n'-1} \kappa^{(s)}_{r+ki} E_{r+ki,r+ki+1}, \quad \kappa^{(s)}_{r+ki} = \frac{\sigma_{r+1}(q) - \sigma_r(q)}{E^{(s)}_{r+ki}},$$

where $(E_{a,b})_{c,d} = \delta_{a,c}\delta_{b,d}$ for $a, b, c, d \in \mathbb{Z}/n\mathbb{Z}$, and $\mathbb{I}_n$ denotes the identity matrix.

Lemma 7.5. (i) *The action of* s_r $(r = 1, \ldots, m-1)$ *on* $\mathcal{M}$ *induces the identity map on* $\mathcal{L}$*, and the action of* π *on* $\mathcal{M}$ *induces the adjoint transformation*

$$\pi^* : L(x) \mapsto Q_1^{-1} L(x) Q_1$$

on $\mathcal{L}$*.*

(ii) *The action of* $\tilde{s}_r$ *and* $\tilde{\pi}$ *on* $\mathcal{M}$ *induces adjoint transformations on* $\mathcal{L}$*, given by*

$$\tilde{s}_r^* : L(x) \mapsto (B_{0,r})^{-1} L(x) B_{0,r},$$

for $r = 0, \ldots, N-1$*, and*

$$\tilde{\pi}^* : L(x) \mapsto P(x)L(x)P(x)^{-1}.$$

Proof. (i) Due to Lemma 7.1, we have $\alpha \circ s_r(Q) = \alpha(Q)$ for $1 \le r \le m-1$. The induced action of π is

$$\pi^*(Q_1 \cdots Q_m P(x)^k) = Q_2 \cdots Q_m P(x)^k Q_1,$$

and the claim follows.

(ii) By [LP13, Theorem 6.2], the action of $\tilde{s}_r$ on $\mathcal{M}$ is given by

$$(Q_s)_{s=1,\ldots,m} \mapsto ((B_{s,r})^{-1} Q_s\, B_{s+1,r})_{s=1,\ldots,m}. \tag{7.11}$$

By definition $E^{(s)}_{r+ki}$ satisfies $E^{(s+m)}_{r+ki} = E^{(s)}_{r+k(i-1)}$, and we have $B_{s+m,r} = P(x)^k B_{s,r} P(x)^{-k}$. The claim follows.

The second part is easy to see, as the transformation $\tilde{\pi}$ just cycles the indices inside the Q_i (Definition 2.3). □

Since similar matrices have the same characteristic polynomial, we obtain the following:

Corollary 7.6. *For each $q \in \mathcal{M}$, $\psi(q)$ is invariant under the action of $W \times \tilde{W}$. In particular, the coefficients of $f(x,y) = \psi(q)$ are conserved quantities for $W \times \tilde{W}$.*

Lemma 7.7. *Each $c \in \mathcal{S}_f$ is invariant under actions of s_i and $\tilde{s}_i$. The cyclic shift π acts on $\mathcal{S}_f$ via*

$$\tau^{-1} : (c_1, \dots, c_M) \mapsto (c_2, \dots, c_M, c_1),$$

and $\tilde{\pi}$ acts on $\mathcal{S}_f$ via

$$\tau : (c_1, \dots, c_M) \mapsto (c_M, c_1, \dots, c_{M-1}).$$

Proof. From the definition of the cyclic group action (2.7), it follows that $\tilde{\pi}(Q)_{ij} = q_{i,j-1}$ and $\pi(Q)_{ij} = q_{i+1,j}$. Thus, using Lemma 6.5, we see that $\tilde{\pi}$ induces

$$c_\ell = \sum_{i+j \equiv \ell+1 \mod M} q_{i,j} \ \mapsto \sum_{i+j \equiv \ell+1 \mod M} q_{i,j-1} = \sum_{i+j \equiv \ell \mod M} q_{i,j} = c_{\ell-1},$$

and that π induces $c_\ell \mapsto c_{\ell+1}$ similarly. □

7.4. **Eigenvector change under conjugation.** For a vector $v = (v_1, v_2, \dots, v_n)$ of rational functions on C_f, define

$$D(v) = (\text{common zeroes of } v_i) - (\text{common poles of } v_i) + R + O_n$$

where R is the divisor supported on $S_{P,O} = \{P, O_1, \dots, O_N\}$, which appears in Theorem 5.1. (See Lemma 6.4 for the explicit formula of R.) The following result is immediate.

Lemma 7.8. *Let f be a rational function on C. Then $D((fv_1, \dots, fv_n))$ is linearly equivalent to $D((v_1, \dots, v_n))$.*

Lemma 7.9. *The positive general divisor $\mathcal{D} = \mathcal{D}(L(x))$ of degree g associated to $L(x)$ is equal to $D((\Delta_{1,n}, \dots, \Delta_{n,n}))$.*

Proof. By Theorem 5.1, $\mathcal{D} = D_n$ belongs to the common zeroes. Also by Theorem 5.1, $\bar{D}_1, \dots, \bar{D}_n$ have no common points, so they do not contribute to $D((\Delta_{1,n}, \dots, \Delta_{n,n}))$. □

Lemma 7.10. *Suppose $M = M(x,y)$ is a $n \times n$ matrix whose entries are polynomials in $\mathbb{C}[x,y]$, thought of as rational functions on C_f with poles supported at P. Let $D(M \cdot v) - D(v) = D_+ - D_-$, where D_+ and D_- are positive divisors. Then restricted to $C_f \setminus P$, we have that $(\det M)_0 - D_+$ is a positive divisor. Also, D_- is supported at P.*

Proof. Let $p \in C_f \setminus P$. Then the entries of M are regular at p. We shall show that the multiplicity of p in D_+ is less than the multiplicity of p as a zero in $\det M$. Since $v'_i = \sum_j M_{ij}(x,y)v_j$, and M_{ij} is regular at p, it is clear that $\text{mult}_p D(Mv) \geq \text{mult}_p D(v)$, where mult_p denotes the multiplicity of a divisor at a point p. We also have

$$v_i = \sum_j (M^{-1})_{ij}(x,y)v'_j = \frac{1}{(\det M)(x,y)} \sum_j \pm |M|_{i,j}(x,y)v'_j.$$

Since $|M|_{i,j}(x,y)$ is regular at p, we have $\text{mult}_p D(v) \geq \text{mult}_p(Mv) - \text{mult}_p(\det M)$. Both claims follow. □

Suppose $L'(x) = Q_1^{-1}L(x)Q_1$ and $\mathcal{D}' = \mathcal{D}(L'(x))$. We shall compute $\mathcal{D}' - \mathcal{D}$ up to linear equivalence. Note that $v = (\Delta_{1,n}, \ldots, \Delta_{n,n})^T$ is an eigenvector of $L(x)^T - y$, and $L'(x)^T = Q_1^T L(x)^T (Q_1^{-1})^T$, so an eigenvector of $L'(x)^T - y$ is equal to $v' = Q_1^T v$, where

$$Q_1^T = Q_1^T(x) = \begin{pmatrix} q_{1,1} & 1 & 0 & 0 \\ 0 & q_{1,2} & 1 & 0 \\ 0 & \ddots & \ddots & 1 \\ x & 0 & 0 & q_{1,n} \end{pmatrix}.$$

Extend the vector v into an infinite vector $\tilde{v}$ by $v_{i+n} = xv_i$. As in §3, let $A = (a_{i,j})_{i \in [n], j \in \mathbb{Z}}$ be the infinite "unfolded" version of $L^T(x)$, satisfying $(L^T)_{i,j}(x) = \sum_\ell a_{i,j+\ell n} x^\ell$. (Note that A is a matrix of scalars, but $\tilde{v} = (\ldots, v_{-1}, v_0, v_1, \ldots)^T$ is a matrix of functions.) Then we have

$$A \cdot \tilde{v} = y\tilde{v}. \tag{7.12}$$

Define $w = (v_1, v_2, \ldots, v_{k+m})^T$ and $w' = (v'_1, v'_2, \ldots, v'_{k+m})^T$. We claim that there exists a $(k+m) \times (k+m)$ matrix $M = M(y)$ such that $w' = M(y) \cdot w$. Since $v' = Q_1^T v$, we have $v'_i = q_i v_i + v_{i+1}$ (where q_i is extended periodically if $i \geq n$). We now write v_{k+m+1} in terms of $v_1, \ldots, v_{k+m}$ using (7.12). The matrix A is supported on the $m+1$ diagonals $k, k+1, \ldots, k+m$. The matrix $A - y$ is supported on the $(k+m+1)$ diagonals $0, 1, \ldots, k+m$. Thus (7.12) gives

$$a_{1,k+1}v_{k+1} + \cdots + a_{1,k+m+1}v_{k+m+1} = yv_1. \tag{7.13}$$

Also note that $a_{1,k+m+1} = 1$. So

$$M(y) = \begin{pmatrix} q_{1,1} & 1 & 0 & \cdots & \cdots & 0 & 0 \\ 0 & q_{1,2} & 1 & 0 & 0 & 0 & 0 \\ \vdots & & \ddots & \ddots & & & \\ 0 & \cdots & & q_{1,k+1} & 1 & & \\ \vdots & & & & \ddots & \ddots & \\ 0 & \cdots & & & & q_{k+m-1} & 1 \\ y & 0 & \cdots & -a_{1,k+1} & \cdots & -a_{1,k+m-1} & q_{1,k+m} - a_{1,k+m} \end{pmatrix}.$$

Lemma 7.11. *With the above conventions, we have*

$$\mathcal{D}' \sim \mathcal{D} + A_1 - P.$$

Proof. We shall show that $D(v') - D(v) = A_1 - P$. This suffices by Lemmas 7.8 and 7.9. From Lemma 7.10, we have $D(v') - D(v) = D_+ - D_-$ where $D_\pm$ are positive divisors, with D_+ when restricted to $C_f \setminus P$ supported on $\{(\det Q^T)(x) = 0\}$, and D_- supported at P. Since $D(v')$ is a positive divisor of degree g by the construction of $L'(x)^T$, we have that D_+ and D_- have the same degree.

We now calculate D_-. By Theorem 5.1, we have that $\mathrm{mult}_P(v_i) = C - i$ for some integer C (we use the convention that negative multiplicity is a pole), and this formula still holds for the infinite vector $(\ldots, v_{-1}, v_0, v_1, \ldots)$. Since $v'_i = q_i v_i + v_{i+1}$, we have $\mathrm{mult}_P(v'_i) = C - (i+1)$. Thus $D_- = P$. It follows that D_+ is a single point in $C_f \setminus P$.

We shall show that D_+ must be supported on A_1. By Theorem 5.1, the common zeros of v_i and v_j for any $i \neq j$ are only D_n except for the points in $S_{P,O}$. Thus we have $D(w) = D_n + R'$ where R' is supported on $S_{P,O}$. But $\{(\det Q^T)(x) = 0\}$ does not intersect $S_{P,O}$. Using Lemma 7.10 now applied to $w' = M(y)w$, we see that D_+ must be supported on $\{\det M(y) = 0\}$.

By Lemma 7.12 below, we conclude that D_+ is a multiple of A_1. Since $D(v') - D(v)$ is a divisor of degree 0, we must have

$$D(v') - D(v) = A_1 - P,$$

as required. □

Lemma 7.12. *The intersection of* $(\det Q^T)(x) = 0$ *and* $(\det M(y)) = 0$ *is the single point* A_1*.*

Proof. We check that $\det M(y) = \pm y$. It is clear that

$$\det M(y) = (-1)^{m+1} y + q_{1,1} \cdots q_{1,k} \det \begin{pmatrix} q_{1,k+1} & 1 & & & \\ & q_{1,k+2} & 1 & & \\ & & \ddots & \ddots & \\ & & & q_{1,k+m-1} & 1 \\ -a_{1,k+1} & -a_{1,k+2} & \cdots & -a_{1,k+m-1} & q_{1,k+m} - a_{1,k+m} \end{pmatrix}.$$

Write u_i $(i = 1, \ldots, m)$ for the m columns of the above m by m matrix. We show that they are linearly dependent as

$$u_1 - q_{1,k+1}\big(u_2 - q_{1,k+2}(u_3 - \cdots (u_{m-1} - q_{1,k+m-1} u_m) \cdots)\big) = 0,$$

which reduces to

$$\begin{aligned} &- a_{1,k+1} + q_{k+1} a_{1,k+2} - q_{1,k+1} q_{1,k+2} a_{1,k+3} + \cdots \\ &+ (-1)^m q_{k+1} q_{k+2} \cdots q_{k+m-1} a_{1,k+m} + (-1)^{m+1} q_{k+1} q_{k+2} \cdots q_{k+m} = 0. \end{aligned} \tag{7.14}$$

By the definition of A, we have

$$a_{1,j} = \begin{cases} l_{j,1} & j = k+1, \ldots, n, \\ l_{j-n,1} & j = n+1, \ldots, n+k, \end{cases}$$

where $L = (l_{ij})_{i \in [n], j \in \mathbb{Z}}$ is the infinite unfolded version of $L(x)$. On the network G', $l_{k+i,1}$ is the weight generating function of paths from the source $k+i$ to the sink $k+1$. Note that each weight in $l_{k+i,1}$ is of degree $m+1-i$ in q-s. We divide $l_{k+i,1}$ into two parts:

$$l_{k+i,1} = l'_{k+i,1} + l''_{k+i,1},$$

where $l'_{k+i,1}$ is the sum of the weights *with* $q_{1,k+i}$, and $l'_{k+i,1}$ is the sum of the weights *without* $q_{1,k+i}$. We claim that there is one-to-one correspondence between the paths contributing to $l'_{k+i,1}$ and the paths contributing to $l''_{k+i+1,1}$. Precisely, we have $q_{1,k+i} l''_{k+i+1,1} = l'_{k+i,1}$. Combining with $l''_{k+1,1} = 0$ and $l'_{k+m,1} = q_{1,k+m}$, we obtain (7.14). □

7.5. **Proof of Theorem 7.2.** (i) Due to the definition of s_r, Lemma 7.5(i) and Lemma 7.7, for $Q = (Q_1, \ldots, Q_m) \in \psi^{-1}(f)$ with $\phi(Q) = ([\mathcal{D}], c, O, (A_1, \ldots, A_m))$, we have

$$\phi \circ s_r(Q_1, \ldots, Q_m) = ([\mathcal{D}], c, O, (A_1, \ldots, A_{r-1}, A_r(Q'), A_{r+1}(Q'), A_{r+2}, \ldots, A_m)),$$

where $Q' = s_r(Q)$. From (2.1) we see that $\epsilon_r(q') = \epsilon_{r+1}(q)$ and $\epsilon_{r+1}(q') = \epsilon_r(q)$, and (7.2) follows.

It is easy to see that for $L(x) = \alpha(Q)$ we have $\eta \circ \pi(L(x)) = ([\mathcal{D}'], \tau^{-1}(c), O')$, where $\mathcal{D}'$ is some divisor. By Lemma 7.11, we have $[\mathcal{D}] = [\mathcal{D}' + A_1 - P]$. The claim follows.

(ii) First we prove (7.4). Let $g = (g_1, \dots, g_n)^t$ be the eigenvector of $L(x) \in \psi^{-1}(f)$, and set $(g'_1, \dots, g'_n)^t := (B_{0,r})^{-1} g$. Define $h'_i := g'_i / g'_n$ for $i = 1, \dots, n-1$. Then we have

$$g'_j = \begin{cases} g_{r+ki} - \kappa^{(0)}_{r+ki} g_{r+ki+1} & j \equiv r + ki \mod n \\ g_j & \text{otherwise.} \end{cases}$$

We must show that $(h'_j)_\infty = (h_j)_\infty$ for $r = 1, \dots, N-1$ and $j = 1, \dots, n-1$. It is enough to consider the case of $r = 1$ and $j = 1$. Due to Theorem 5.1 and (5.1), there are positive divisors $\mathcal{D}, \mathcal{D}'$ of degree g such that $(h_1)_\infty = (n-1)P + \mathcal{D}$, $(h'_1)_\infty = (n-1)P + \mathcal{D}'$. Since $h'_1 = h_1 - \kappa^{(0)}_1 h_2$ and $(h_1)_\infty > (\kappa^{(0)}_1 h_2)_\infty = (h_2)_\infty$, we get $(h'_1)_\infty \le (h_1)_\infty$. Thus it follows that $\mathcal{D}' = \mathcal{D}$. From (7.11) and Lemma 7.7 it follows that both $A \in R_A$ and $c \in \mathbb{C}^M$ are not changed by $\tilde{s}_r$. From the facts that $\sigma_{N+1-r}(q)$ is the product of all diagonal elements of $\tilde{Q}_r$, and that the R-matrix action changes $(\tilde{Q}_r, \tilde{Q}_{r+1})$ to $(\tilde{Q}'_{r+1}, \tilde{Q}'_r)$, it follows that $\tilde{s}_r$ exchanges the $(N-r)$-th and the $(N-r+1)$-th elements of $O \in R_O$. Then we get (7.4).

Since $\tilde{\pi}$ induces the shift ς^* of $L(x)$, (7.5) follows from Proposition 7.4.

7.6. **Snake path actions.** Recall that $M = \gcd(n, k+m)$. The *snake path actions* are torus actions on $\mathcal{M}$, considered in [LP13]. Recall from § 4.2 that a snake path is a closed path on G that turns at every vertex. Thus it alternates between going up or going right. For $1 \le s \le M$ and $t \in \mathbb{C}^*$, the action $T_s := T_s(t)$ on $\mathcal{M}$ is given by

$$T_s(q)_{ij} = \begin{cases} t\, q_{ij} & \text{if } j \equiv s - i + 1 \mod M \\ \dfrac{1}{t}\, q_{ij} & \text{if } j \equiv s - i \mod M \\ q_{ij} & \text{otherwise.} \end{cases} \tag{7.15}$$

Informally, $T_s(t)$ multiplies all left turns on a snake path by t, and all right turns by $1/t$.

Lemma 7.13. *The induced action T^*_s $(1 \le s \le M)$ on $\mathcal{L}$ is given by*

$$T^*_s : \; L(x) \mapsto D_s(t) L(x) D_s(t)^{-1},$$

where

$$D_s(t) = \mathrm{diag}(d_i)_{1 \le i \le n}, \quad d_i = \begin{cases} t & \text{if } i \equiv s \mod M \\ 1 & \text{otherwise}. \end{cases}$$

Proof. For simplicity we write $D_s := D_s(t)$ and $P := P(x)$. We rewrite $D_s L(x) D_s^{-1}$ as

$$D_s L(x) D_s^{-1} = \left(\prod_{i=1}^{m} (P^{-i+1} D_s P^{i-1}) Q_i (P^{-i} D_s P^i)^{-1} \right) \cdot (P^{-m} D_s P^m) P^k D_s^{-1}. \tag{7.16}$$

This is equal to $\alpha(T_s(Q))$ since $(P^{-m} D_s P^m) \cdot P^k D_s^{-1} = P^k$. □

Proposition 7.14. *We have the following commutative diagram:*

$$\begin{tikzcd} \mathcal{M} \arrow[r, "T_s"] \arrow[d, "e_u"'] & \mathcal{M} \arrow[d, "e_u"] \\ \mathcal{M} \arrow[r, "\varsigma^{-u} \circ T_s \circ \varsigma^u"'] & \mathcal{M}. \end{tikzcd}$$

Proof. It is enough to prove $((\varsigma^*)^{-u}\circ T_s^*\circ(\varsigma^*)^u)\circ e_u^* = e_u^*\circ T_s^*$ acting on $\mathcal{L}$. For $L(x) = Q_1\cdots Q_m P^k \in \mathcal{L}$, we have

$$\begin{aligned}
L(x) &\overset{e_u^*}{\longmapsto} Q_{u+1}\cdots Q_m P^k Q_1\cdots Q_u \\
&\overset{(\varsigma^*)^{-u}\circ\tilde{T}_s\circ(\varsigma^*)^u}{\longmapsto} (P^{-u}D_sP^u)Q_{u+1}\cdots Q_mP^kQ_1\cdots Q_u(P^{-u}D_sP^u)^{-1} \\
&= \prod_{i=u+1}^{m}(P^{-i+1}D_sP^{i-1})Q_i(P^{-i}D_sP^i)^{-1}\cdot(P^{-m}D_sP^m)P^kD_s^{-1} \\
&\qquad\qquad \cdot\prod_{i=1}^{u}(P^{-i+1}D_sP^{i-1})Q_i(P^{-i}D_sP^i)^{-1} \\
&= Q'_{u+1}\cdots Q'_mP^kQ'_1\cdots Q'_u,
\end{aligned}$$

where $Q'_i = (P^{-i+1}D_sP^{i-1})Q_i(P^{-i}D_sP^i)^{-1}$. On the other hand we have

$$\begin{aligned}
L(x) &\overset{T_s^*}{\longmapsto} D_sL(x)D_s^{-1} \overset{(7.16)}{=} Q'_1\cdots Q'_mP^k \\
&\overset{e_u^*}{\longmapsto} Q'_{u+1}\cdots Q'_mP^kQ'_1\cdots Q'_u.
\end{aligned}$$

Thus the claim is obtained. $\square$

From Lemma 6.5 and 7.13 we obtain

Proposition 7.15. *The following diagram is commutative:*

$$\begin{tikzcd}
\psi^{-1}(f) \arrow[r, "\phi"] \arrow[d, "T_s"'] & \mathrm{Pic}^g(C_f)\times\mathcal{S}_f\times R_O\times R_A \arrow[d, "{(id, t_s, id, id)}"] \\
\psi^{-1}(f) \arrow[r, "\phi"'] & \mathrm{Pic}^g(C_f)\times\mathcal{S}_f\times R_O\times R_A,
\end{tikzcd} \tag{7.17}$$

where $t_s := t_s(t)$ acts on $\mathcal{S}_f$ by

$$t_s(c_\ell) = \begin{cases} t\,c_\ell & \textit{if } s=\ell \\ \dfrac{1}{t}\,c_\ell & \textit{if } s=\ell+1 \\ c_\ell & \textit{otherwise.}\end{cases} \tag{7.18}$$

Thus the snake path actions $T_s(t)$ act transitively on $\mathcal{S}_f$.

8. THETA FUNCTION SOLUTION TO INITIAL VALUE PROBLEM

8.1. **Riemann theta function.** Fix $f\in\mathring{\mathcal{V}}$. We fix a universal cover of C_f and $P_0\in C_f$, and define the Abel-Jacobi map ι by

$$\iota : C_f\to\mathbb{C}^g;\ X\mapsto\left(\int_{P_0}^X\omega_1,\dots,\int_{P_0}^X\omega_g\right),$$

where $\omega_1,\dots,\omega_g$ is a basis of holomorphic differentials on C_f. We also write ι for the induced map $\mathrm{Div}^0(C_f)\to\mathbb{C}^g$. Let Ω be the period matrix of C_f, and write $\Theta(z) := \Theta(z;\Omega)$ for the Riemann theta function:

$$\Theta(z;\Omega) = \sum_{\mathbf{m}\in\mathbb{Z}^g}\exp\left(\pi\sqrt{-1}\mathbf{m}\cdot(\Omega\mathbf{m}+2z)\right),\quad z\in\mathbb{C}^g.$$

It is known to satisfy the quasi-periodicity:

$$\Theta(z+\mathbf{m}+\mathbf{n}\Omega)=\exp\left(-\pi\sqrt{-1}\,\mathbf{n}\cdot(\Omega\mathbf{n}+2z)\right)\Theta(z),$$

for $\mathbf{m},\mathbf{n}\in\mathbb{Z}^g$.

Let $\mathcal{D}_0\in\mathrm{Div}^{g-1}(C_f)$ be the Riemann characteristic, that is, $2\mathcal{D}_0$ is linearly equivalent to the canonical divisor K_{C_f}. Then the well known properties of the Riemann theta function gives

Lemma 8.1. *For any point $Y\in C_f$, the function*

$$X\mapsto\Theta(\iota(X-Y))$$

is a section of the line bundle $\mathcal{O}(\mathcal{D}_0+Y)$ which has zeroes exactly at $\mathcal{D}_0$ and at Y, and no poles. For any positive divisor $\mathcal{D}$ of degree g, the function

$$X\mapsto\Theta(\iota(X-\mathcal{D}+\mathcal{D}_0))$$

is a section of the line bundle $\mathcal{O}(\mathcal{D})$ which has zeroes exactly at $\mathcal{D}$ and no poles.

8.2. **The inverse of ϕ.** For a positive divisor $\mathcal{D}$ of degree g, define the function ψ_i ($i\in\mathbb{Z}$) on C_f by

$$\psi_i(X)=\frac{\Theta(\beta-\iota(\mathcal{O}_{n-i})+\iota(X-O_n))\prod_{\ell=i+1}^{n}\Theta(\iota(X-O_\ell))}{\Theta(\beta+\iota(X-O_n))\Theta(\iota(X-P))^{n-i}},$$

where $\beta=\iota(\mathcal{D}_0+O_n-\mathcal{D})$.

Lemma 8.2. *$\psi_i(X)$ is a meromorphic function on C_f. When $\mathcal{D}$ is the positive divisor defined by Definition 5.2, we have*

$$(\psi_i)=(h_i)=\mathcal{D}_i+\sum_{j=i+1}^{n}O_j-\mathcal{D}-(n-i)P. \tag{8.1}$$

Thus $\psi_i(X)$ is equal to $h_i(X)$ up to a scalar.

Proof. To check that $\psi_i(X)$ is a meromorphic function on C_f we use the functional equation for Θ, and note that the multiset of points (with signs and multiplicities) that appear in the numerator is equal to the same for the denominator. This multiset in additive notation is $\beta-(n-i)P+(n-i+1)X-O_n$. For the second statement, we note that $\psi_i(X)$ has zeroes at $O_{i+1},O_{i+2},\dots,O_n$, a pole of order $n-i$ at P, and also poles at $\mathcal{D}$. By (5.1), we have $(\psi_i)=(h_i)$. □

Now we study the inverse of ϕ. We focus on the $\mathbb{Z}^m$ action. For $t=(t_1,t_2,\dots,t_m)\in\mathbb{Z}^m$, let $q^t:=(q^t_{ji})_{ji}\in\psi^{-1}(f)$ be a configuration at time t. For $L^t(x):=\alpha(q^t)$, let $g^t=(g^t_1,\dots,g^t_n)^\perp$ be its eigenvector and set $h^t_i:=g^t_i/g^t_n$. Let $\phi(q^t)$ be

$$(\mathcal{D}^t,(c^t_1,\cdots,c^t_M),(A_1,\dots,A_m),(O_1,\dots,O_N)),$$

where $\mathcal{D}^t=\mathcal{D}-\sum_{j=1}^{m}t_j\mathcal{A}_j$ for some positive divisor $\mathcal{D}$ of degree g. Define $\mathbf{e}_j:=(\underbrace{1,\dots,1}_{j},\underbrace{0,\dots,0}_{m-j})$ for $j=0,\dots,m-1$, and recursively set $\mathbf{e}_{j+m}=(1,\dots,1)+\mathbf{e}_j$. Define functions θ^t_i and Ψ^t_i on C_f by

$$\theta^t_i(X)=\Theta(\beta^t+\iota(X-O_n-\mathcal{O}_{n-i})),$$

where $\beta^t = \iota(\mathcal{D}_0 + O_n - \mathcal{D}^t)$, and by

$$\Psi_i^{t+\mathbf{e}_j}(X) = \frac{\theta_{i-N}^{t+\mathbf{e}_{j-1}}(X)\, \theta_i^{t+\mathbf{e}_j}(X)}{\theta_i^{t+\mathbf{e}_{j-1}}(X)\, \theta_{i-N}^{t+\mathbf{e}_j}(X)}.$$

By Lemma 8.2, there is a constant $b_i^{t+\mathbf{e}_j}$ depending on C_f, such that we have

$$\Psi_i^{t+\mathbf{e}_j}(X) = b_i^{t+\mathbf{e}_j} \frac{h_{i-N}^{t+\mathbf{e}_{j-1}}(X)\, h_i^{t+\mathbf{e}_j}(X)}{h_i^{t+\mathbf{e}_{j-1}}(X)\, h_{i-N}^{t+\mathbf{e}_j}(X)}.$$

Lemma 8.3. *(Cf. Lemma 3.1 [I10]) We have*

$$\Psi_{i-1}^{t+\mathbf{e}_j}(P) = \Psi_i^{t+\mathbf{e}_j}(O_i) = b_i^{t+\mathbf{e}_j} \frac{q_{j,i-N}^t}{q_{j,i}^t}, \tag{8.2}$$

$$\Psi_i^{t+\mathbf{e}_j}(P) = b_i^{t+\mathbf{e}_j} \frac{d_{i-N-1}^{t+\mathbf{e}_j}\, d_i^{t+\mathbf{e}_j}}{d_{i-1}^{t+\mathbf{e}_j}\, d_{i-N}^{t+\mathbf{e}_j}}, \tag{8.3}$$

where d_i^t is a coefficient of the leading term of $h_i^t(X)$ around $X = P$.

Proof. The first equality of (8.2) follows from $\theta_i^t(O_i) = \theta_{i-1}^t(P)$ and $O_i = O_{i-N}$. By Lemma 7.5 we have $g^{t+\mathbf{e}_{j-1}} = Q_j^t\, g^{t+\mathbf{e}_j}$ which gives

$$\Psi_i^{t+\mathbf{e}_j}(X) = b_i^{t+\mathbf{e}_j} \frac{(h_{i-N-1}^{t+\mathbf{e}_j}(X) + q_{j,i-N}\, h_{i-N}^{t+\mathbf{e}_j}(X))h_i^{t+\mathbf{e}_j}(X)}{(h_{i-1}^{t+\mathbf{e}_j}(X) + q_{j,i}\, h_i^{t+\mathbf{e}_j}(X))h_{i-N}^{t+\mathbf{e}_j}(X)}.$$

Then, from (8.1) the second equalities of (8.2) and (8.3) follow. □

Lemma 8.4. *We have*

$$\frac{d_i^{t+\mathbf{e}_j}}{d_{i+1}^{t+\mathbf{e}_j}} = c_{i+j}^t, \qquad i + j \equiv \ell \mod M, \tag{8.4}$$

where we extend c_i^t to $i \in \mathbb{Z}$ by setting $c_i^t := c_\ell^t$ if $i \equiv \ell \mod M$ for $1 \le \ell \le M$.

Proof. It is enough to show the case of $j = 0$, $1 \le i \le M$. Then the other cases follow from Lemma 7.7.

We will show that the coefficient of the leading term in $\frac{g_i}{g_{i+1}}$ at $X = P$ is equal to c_i. To obtain the ratio $\frac{g_i}{g_{i+1}}$ we may choose to compute maximal minors of $L(x) - y$ with respect to any row. Let us choose row $i + 1$. Then by Theorem 6.2, we are counting families of paths that do not start at source $i + 1$ and do not end at sink i (for g_i) or sink $i + 1$ (for g_{i+1}).

Since we are computing at $X = P$, the only terms that contribute are the ones lying on the upper hull of the Newton polygon, that is, the edge with slope $-n/(m+k)$. In terms of path families this means that we only take paths that are as close to a snake path as possible. In the case of g_{i+1} this means that we take a subset of closed snake paths. Each of them picks up no weight, and thus overall we just get a constant. In the case of g_i one of the closed staircase paths skips one term, and thus we get a long path starting at source i, winding around the torus and ending at sink $i + 1$. Such path essentially by definition picks up weight c_i. The other snake paths may or may not appear, thus creating only a constant factor in front. □

Note that (8.4) is compatible with the snake path actions in Lemma 7.13.

Theorem 8.5. *When $N = 1$, the inverse map of ϕ is given by*

$$q^t_{j,i} = f_c^{-\frac{1}{M}} \cdot a_j \cdot c^t_{i+j-1} \cdot \frac{\theta_i^{t+\mathbf{e}_j}(P)\,\theta_{i-1}^{t+\mathbf{e}_{j-1}}(P)}{\theta_i^{t+\mathbf{e}_{j-1}}(P)\,\theta_{i-1}^{t+\mathbf{e}_j}(P)},$$

where

$$a_j = x(A_j)^{\frac{1}{n}} \cdot \exp\left[\frac{2\pi\sqrt{-1}}{n} b \cdot \iota(P - A_j)\right], \tag{8.5}$$

and $b \in \mathbb{Z}^g$ is determined by $a, b \in \mathbb{Z}^g$ such that $a + b\Omega := \iota(\mathcal{O}_n)$.

Proof. Let $p^t_{j,i} = q^t_{j,i} q^t_{j,i-1} \cdots q^t_{j,i-N+1}$. By Lemma 8.3 we get the equations

$$\alpha^t_j := \frac{p^t_{j,i}}{\Psi_i^{t+\mathbf{e}_j}(P)} \frac{d_i^{t+\mathbf{e}_j}}{d_{i-N}^{t+\mathbf{e}_j}} = \frac{p^t_{j,i-1}}{\Psi_{i-1}^{t+\mathbf{e}_j}(P)} \frac{d_{i-1}^{t+\mathbf{e}_j}}{d_{i-N-1}^{t+\mathbf{e}_j}} = \cdots .$$

We show that α^t_j does not depend on t. Consider the product

$$x(A_j) = p^t_{j,i} p^t_{j,i+N} \cdots p^t_{j,i+(n'-1)N},$$

which is equal to

$$(\alpha^t_j)^{n'} \frac{d_{i-N}^{t+\mathbf{e}_j}}{d_{i+n-N}^{t+\mathbf{e}_j}} \frac{\theta_{i-N}^{t+\mathbf{e}_{j-1}}(P)\,\theta_{i+n-N}^{t+\mathbf{e}_j}(P)}{\theta_{i+n-N}^{t+\mathbf{e}_{j-1}}(P)\,\theta_{i-N}^{t+\mathbf{e}_j}(P)}.$$

Since $\mathcal{O}_n$ is equivalent to 0 in $\mathrm{Pic}^0(C_f)$, there exists $a, b \in \mathbb{Z}^g$ such that $a + b\Omega = \iota(\mathcal{O}_n)$. Due to the quasi-periodicity of $\Theta(z)$, we have

$$\theta^t_{i+n}(P) = \exp\left[-2\pi\sqrt{-1} b \cdot \left(\beta^t - \iota(\mathcal{O}_{n-i} - \mathcal{O}_1) + b\Omega/2\right)\right] \cdot \theta^t_i(P). \tag{8.6}$$

By using (8.4) and (8.6) we obtain α^t_j independent of t as

$$\alpha^t_j = f_c^{-\frac{N}{M}} a_j^N.$$

Hence we get

$$p^t_{j,i} = f_c^{-\frac{N}{M}} a_j^N \cdot \prod_{\ell=1}^N c^t_{i+j-\ell} \cdot \frac{\theta_i^{t+\mathbf{e}_j}(P)\,\theta_{i-N}^{t+\mathbf{e}_{j-1}}(P)}{\theta_i^{t+\mathbf{e}_{j-1}}(P)\,\theta_{i-N}^{t+\mathbf{e}_j}(P)}. \tag{8.7}$$

When $N = 1$, it is nothing but the claim. □

8.3. **Conditional solution for $N > 1$.** When $N > 1$, by factorizing (8.7) we obtain

$$q^t_{j,i} = f_c^{-\frac{1}{M}} \cdot \gamma_{j,i} \cdot a_j \cdot c^t_{i+j-1} \frac{\theta_i^{t+\mathbf{e}_j}(P)\,\theta_{i-1}^{t+\mathbf{e}_{j-1}}(P)}{\theta_i^{t+\mathbf{e}_{j-1}}(P)\,\theta_{i-1}^{t+\mathbf{e}_j}(P)}. \tag{8.8}$$

Here $\gamma_{j,i}$ satisfies

$$\prod_{\ell=1}^N \gamma_{j,i+\ell} = 1, \tag{8.9}$$

which denotes that $\gamma_{j,i} = \gamma_{j,i+N}$.

Lemma 8.6. *For i such that $i \equiv r \mod N$ we have*

$$\prod_{j=1}^m \gamma_{j,i} = \sigma_r^{\frac{1}{n'}} \cdot \prod_{j=1}^m a_j^{-1} \cdot \exp\left(2\pi\sqrt{-1}(b' - b\frac{k'}{n'}) \cdot \iota(P - O_r)\right). \tag{8.10}$$

Here b' is given by $a', b' \in \mathbb{Z}^g$ such that $a' + b'\Omega := \iota(\mathcal{A}_m + \mathcal{O}_k)$.

Proof. From (3.6) and (8.8) it follows that

$$\sigma_r = \left(\prod_{j=1}^{m} \gamma_{j,i}\, a_j\, f_c^{-\frac{1}{M}}\right)^{n'} \cdot \prod_{j=1}^{m}\prod_{p=0}^{n'-1} c^t_{j+i+pN-1} \cdot \prod_{j=0}^{n'-1} \frac{\theta^t_{i+jN-1}(P)\,\theta^{t+\mathbf{e}_m}_{i+jN}(P)}{\theta^{t+\mathbf{e}_m}_{i+jN-1}(P)\,\theta^t_{i+jN}(P)}, \tag{8.11}$$

for $r = 1, \dots, N$ and $i \equiv r \mod N$.

First we show that the second factor of (8.11) is equal to $f_c^{\frac{mn'}{M}}$. It is enough to prove in the case of $i = 1$, where the second factor is $\prod_{p=0}^{n'-1}(c_{1+pN}\cdots c_{m+pN})$. When $n' = k'$, M is a divisor of m and we have $c_{1+pN}\cdots c_{m+pN} = f_c^{m/M}$. When $n' \neq k'$, define an automorphism ν of $\mathcal{N}' := \{0, \dots, n'-1\}$ by $\nu : p \mapsto p+n'-k' \mod n'$. Since n' and k' are coprime, a set $\{\nu^\ell(1) \mid \ell \in \mathcal{N}'\}$ coincides with $\mathcal{N}'$. Thus, from $M = \gcd(n, k+m)$ and $m+pN \equiv \nu(p)N \mod M$, it follows that the product $\prod_{p=0}^{n'-1}(c_{1+pN}\cdots c_{m+pN})$ is reordered to be $\prod_{j=1}^{mn'} c_j$. Further, since M is a divisor of mn', we obtain the claim. Next, we can show that the third factor of (8.11) is equal to $\exp\left(2\pi\sqrt{-1}(-b'n' + bk') \cdot \iota(P - O_r)\right)$, by using (8.6) and

$$\theta_i^{t+\mathbf{e}_m}(P) = \exp\left[-2\pi\sqrt{-1}b' \cdot \left(\beta^t - \iota(\mathcal{O}_{n-i+k} - \mathcal{O}_1) + b'\Omega/2\right)\right] \cdot \theta^t_{i-k}(P).$$

Here b' is given by $a', b' \in \mathbb{Z}^g$ such that $a' + b'\Omega := \iota(\mathcal{A}_m + \mathcal{O}_k)$. Finally from (8.11) we obtain

$$\left(\prod_{j=1}^{m} \gamma_{j,i}\right)^{n'} = \sigma_r \cdot \prod_{j=1}^{m} a_j^{-n'} \cdot \exp\left(2\pi\sqrt{-1}(n'b' - bk') \cdot \iota(P - O_r)\right),$$

and the claim follows. □

Unfortunately, we have not been able to obtain $\gamma_{j,i}$ from (8.9) and (8.10). Nevertheless, if we assume that $\gamma_{j,i}$ is constant on $\mathrm{Pic}^g(C_f)$, then we obtain the following conditional result.

Proposition 8.7. *Suppose that $\gamma_{j,i}$ is a constant function on $\mathrm{Pic}^g(C_f)$. Then we have*

$$\gamma_{j,i} = \sigma_r^{\frac{1}{n'm}} \cdot \prod_{j=1}^{m} a_j^{-\frac{1}{m}} \cdot \exp\left(\frac{2\pi\sqrt{-1}}{n'm}(n'b' - k'b) \cdot \iota(P - O_r)\right). \tag{8.12}$$

In particular, the inverse of ϕ is given by

$$q^t_{j,i} = Q \cdot a_j \cdot o_i \cdot c^t_{i+j-1} \frac{\theta_i^{t+\mathbf{e}_j}(P)\,\theta_{i-1}^{t+\mathbf{e}_{j-1}}(P)}{\theta_i^{t+\mathbf{e}_{j-1}}(P)\,\theta_{i-1}^{t+\mathbf{e}_j}(P)},$$

where a_j is defined by (8.5), *and*

$$Q = f_c^{-\frac{1}{M}} \cdot \prod_{j=1}^{m} x(A_j)^{-\frac{1}{nm}},$$

$$o_i = \left((-1)^{n'+1}\frac{y(O_r)^{n'}}{x(O_r)^{k'}}\right)^{\frac{1}{n'm}} \cdot \exp\left[\frac{2\pi\sqrt{-1}}{nm}\left((nb' - kb) \cdot \iota(P - O_i) - b \cdot \iota(\mathcal{A}_m)\right)\right],$$

$$i \equiv r \mod N.$$

Proof. To have (8.8) compatible with the snake path action (7.15) and (7.18), $\gamma_{i,j}$ has to be constant on $\mathcal{S}_f$. So $\gamma_{j,i}$ is regarded as a function on $R_A \times R_O$. If $\gamma_{j,i}$ is not constant on R_A, (8.10) implies that $\gamma_{j,i}$ depends on a_j^{-1}, but this contradicts (8.9). Thus $\gamma_{j,i}$ is constant on R_A, and we have (8.12) which fulfills (8.9). By substituting (8.12) in (8.8) and using (3.11), we obtain the final claim. □

9. OCTAHEDRON RECURRENCE

We shall prove that the function $\theta_i^t(P)$ satisfies a family of octahedron recurrences [Spe], as a specialization of Fay's trisecant identity. We follow the definitions in §8. In the following we assume $N = \gcd(n,k) = 1$ and write O for the unique special point O_1 over $(0,0) \in \tilde{C}_f$.

9.1. **Fay's trisecant identity.** We introduce Fay's trisecant identity in our setting. For α, $\beta \in \mathbb{R}^g$, we define a generalization of the Riemann theta function:

$$\Theta[\alpha,\beta](z) := \exp\big(\pi\sqrt{-1}\beta\cdot(\beta\Omega + 2z + 2\alpha)\big)\cdot\Theta(z + \Omega\beta + \alpha). \tag{9.1}$$

We call $[\alpha,\beta]$ *a half period* when α, $\beta \in (\mathbb{Z}/2)^g$. Furthermore, a half period $[\alpha,\beta]$ is *odd* when $\Theta[\alpha,\beta](z)$ is an odd function of z, that is, $\Theta[\alpha,\beta](-z) = -\Theta[\alpha,\beta](z)$. We note that the Riemann theta function itself is an even function: $\Theta(z) = \Theta(-z)$. It is easy to check that a half period $[\alpha,\beta]$ is odd if and only if $4\alpha\cdot\beta \equiv 1 \mod 2$.

Theorem 9.1. *[Fay, §II] For four points P_1, P_2, P_3, P_4 on the universal cover of C_f, $z \in \mathbb{C}^g$, and an odd half period $[\alpha,\beta]$, the formula*

$$\begin{aligned}
&\Theta(z+\iota(P_3-P_1))\,\Theta(z+\iota(P_4-P_2))\,\Theta[\alpha,\beta](\iota(P_2-P_3))\,\Theta[\alpha,\beta](\iota(P_4-P_1))\\
&+\Theta(z+\iota(P_3-P_2))\,\Theta(z+\iota(P_4-P_1))\,\Theta[\alpha,\beta](\iota(P_1-P_3))\,\Theta[\alpha,\beta](\iota(P_2-P_4))\\
&=\Theta(z+\iota(P_3+P_4-P_1-P_2))\,\Theta(\mathbf{z})\,\Theta[\alpha,\beta](\iota(P_4-P_3))\,\Theta[\alpha,\beta](\iota(P_2-P_1)).
\end{aligned}$$

holds.

9.2. $m=2$ **case.** When $m=2$, the vertical actions e_u $(u=1,2)$ are written as difference equations expressed as

$$q_{2,i}^t q_{1,i-k}^t = q_{1,i}^{t+\mathbf{e}_1} q_{2,i}^{t+\mathbf{e}_1}, \tag{9.2}$$

$$q_{2,i+1}^t + q_{1,i-k}^t = q_{1,i+1}^{t+\mathbf{e}_1} + q_{2,i}^{t+\mathbf{e}_1}, \tag{9.3}$$

$$q_{j,i-k}^t = q_{j,i}^{t+\mathbf{e}_2} \qquad (j=1,2). \tag{9.4}$$

For simplicity we write θ_i^t for $\theta_i^t(P) = \Theta(\beta^t + \iota((n-i-1)(O-P)))$. By construction the theta function solution of $q_{j,i}^t$ (Theorem 8.5),

$$q_{j,i}^t = f_c^{-\frac{1}{M}}\cdot a_j\cdot c_{i+j-1}^t\cdot\frac{\theta_i^{t+\mathbf{e}_j}\,\theta_{i-1}^{t+\mathbf{e}_{j-1}}}{\theta_i^{t+\mathbf{e}_{j-1}}\,\theta_{i-1}^{t+\mathbf{e}_j}}, \tag{9.5}$$

satisfies (9.2)–(9.4).

Theorem 9.2. *For any $t \in \mathbb{Z}^2$ and $i \in \mathbb{Z}$, the θ_i^t satisfy an octahedron recurrence relation,*

$$a_2\,\theta_{i+1}^{t+\mathbf{e}_2}\theta_i^{t+2\mathbf{e}_1} - a_1\,\theta_{i+1}^{t+2\mathbf{e}_1}\theta_i^{t+\mathbf{e}_2} = c\,\theta_i^{t+\mathbf{e}_1+\mathbf{e}_2}\theta_{i+1}^{t+\mathbf{e}_1}. \tag{9.6}$$

Here c is a constant given by

$$c = a_2\,\frac{\Theta(p_0+\iota(A_1-A_2))\Theta(p_0+\iota(O-P))}{\Theta(p_0+\iota(O-A_2))\Theta(p_0+\iota(A_1-P))},$$

where p_0 is a zero of the Riemann theta function: $\Theta(p_0)=0$.

Proof. By setting $(P_1, P_2, P_3, P_4) = (A_2, O, P, A_1)$ and using (9.1), we obtain:

$$\begin{aligned} &T_1\, \Theta(z + \iota(P - A_2))\, \Theta(z + \iota(A_1 - O)) + T_2\, \Theta(z + \iota(P - O))\, \Theta(z + \iota(A_1 - A_2)) \\ &\qquad = T_3\, \Theta(z + \iota(P + A_1 - A_2 - O))\, \Theta(z) \end{aligned} \tag{9.7}$$

where $p := \Omega\beta + \alpha$, and

$$\begin{aligned} T_1 &:= \Theta(p + \iota(O - P))\, \Theta(p + \iota(A_1 - A_2)), \\ T_2 &:= \mathrm{e}^{4\pi\sqrt{-1}\beta\cdot\iota(A_2 - A_1)}\, \Theta(p + \iota(A_2 - P))\, \Theta(p + \iota(O - A_1)), \\ T_3 &:= \Theta(p + \iota(A_1 - P))\, \Theta(p + \iota(O - A_2)). \end{aligned} \tag{9.8}$$

By setting $z = \beta^t + \iota((n - i - 1)(O - P) + 2\mathcal{A}_1)$, (9.7) turns out to be

$$T_1\, \theta_i^{\mathbf{e}_1 + \mathbf{e}_2}\, \theta_{i+1}^{\mathbf{e}_1} + T_2\, \theta_{i+1}^{2\mathbf{e}_1}\, \theta_i^{\mathbf{e}_2} = T_3\, \theta_{i+1}^{\mathbf{e}_2}\, \theta_i^{2\mathbf{e}_1}. \tag{9.9}$$

Lemma 9.3. *We have*

$$\frac{T_2}{T_3} = \frac{a_1}{a_2}.$$

Proof. For $q^0 \in \psi^{-1}(f)$, take $t_0 \in \mathbb{Z}^m$ such that $\beta^{t_0} \in \mathbb{C}^g$ is a zero of the Riemann theta function, which is always possible by choosing q^0 appropriately. We write $-p_0$ for such β^{t_0}. Then we have $\theta_{n-1}^{t_0} = \Theta(-p_0) = 0$, and $q_{1,n}^{t_0} = q_{2,n}^{t_0 - \mathbf{e}_1} = 0$. From (9.2) and (9.3), we obtain

$$q_{2,n-1}^{t_0 - \mathbf{e}_1} q_{1,n-1-k}^{t_0 - \mathbf{e}_1} = q_{1,n-1}^{t_0} q_{2,n-1}^{t_0}, \qquad q_{1,n-k-1}^{t_0 - \mathbf{e}_1} = q_{2,n-1}^{t_0}. \tag{9.10}$$

On the other hand, when $z = p_0 + \iota(A_2 - P)$, (9.7) becomes

$$T_2\, \Theta(p_0 + \iota(A_2 - O))\, \Theta(p_0 + \iota(A_1 - P)) = T_3\, \Theta(p_0 + \iota(A_1 - O))\, \Theta(p_0 + \iota(A_2 - P)).$$

It is rewritten as (using that $\Theta(z)$ is even function)

$$\frac{T_2}{T_3} = \frac{\theta_{n-2}^{t_0 + \mathbf{e}_1}\, \theta_{n-1}^{t_0 + \mathbf{e}_2 - \mathbf{e}_1}}{\theta_{n-2}^{t_0 + \mathbf{e}_2 - \mathbf{e}_1}\, \theta_{n-1}^{t_0 + \mathbf{e}_1}} = \frac{a_1}{a_2} \cdot \frac{q_{2,n-1}^{t_0 - \mathbf{e}_1}}{q_{1,n-1}^{t_0}},$$

where we use (9.5) to get the last equality. It follows from (9.10) that $q_{2,n-1}^{t_0 - \mathbf{e}_1} / q_{1,n-1}^{t_0} = 1$, and we obtain the claim. □

We continue the proof of the theorem. By setting $z = p_0 + \iota(O - P)$ in (9.7), we obtain

$$T_1\, \Theta(p_0 + \iota(O - A_2))\, \Theta(p_0 + \iota(A_1 - P)) = T_3\, \Theta(p_0 + \iota(A_1 - A_2))\, \Theta(p_0 + \iota(O - P)).$$

Using this and the above lemma, (9.9) is shown to be

$$c\, \theta_i^{\mathbf{e}_1 + \mathbf{e}_2}\, \theta_{i+1}^{\mathbf{e}_1} + a_1\, \theta_{i+1}^{2\mathbf{e}_1}\, \theta_i^{\mathbf{e}_2} = a_2\, \theta_{i+1}^{\mathbf{e}_2}\, \theta_i^{2\mathbf{e}_1}.$$ □

Conversely, we have the following.

Proposition 9.4. *Suppose that* θ_i^t *satisfy* (9.6). *Then* $q_{j,i}^t$ *defined by* (9.5) *satisfies* (9.2)–(9.4).

Proof. It is very easy to see that (9.5) satisfies (9.2) and (9.4). We check (9.3). We consider a ratio

$$f_i^t := \frac{q_{2,i+1}^t - q_{1,i+1}^{t + \mathbf{e}_1}}{q_{2,i}^{t + \mathbf{e}_1} - q_{1,i}^{t + \mathbf{e}_2}}.$$

By substituting (9.5) in f_i^t we obtain

$$f_i^t = \frac{a_2 c_{i+2} \frac{\theta_{i+1}^{\mathbf{e}_2}\,\theta_i^{\mathbf{e}_1}}{\theta_{i+1}^{\mathbf{e}_1}\,\theta_i^{\mathbf{e}_2}} - a_1 c_{i+1}^{\mathbf{e}_1} \frac{\theta_{i+1}^{2\mathbf{e}_1}\,\theta_i^{\mathbf{e}_1}}{\theta_{i+1}^{\mathbf{e}_1}\,\theta_i^{2\mathbf{e}_1}}}{a_2 c_{i+1}^{\mathbf{e}_1} \frac{\theta_i^{\mathbf{e}_1+\mathbf{e}_2}\,\theta_{i-1}^{2\mathbf{e}_1}}{\theta_i^{2\mathbf{e}_1}\,\theta_{i-1}^{\mathbf{e}_1+\mathbf{e}_2}} - a_1 c_i^{\mathbf{e}_2} \frac{\theta_i^{\mathbf{e}_1+\mathbf{e}_2}\,\theta_{i-1}^{\mathbf{e}_2}}{\theta_i^{\mathbf{e}_2}\,\theta_{i-1}^{\mathbf{e}_1+\mathbf{e}_2}}} = \frac{\theta_i^{\mathbf{e}_1}\,\theta_{i-1}^{\mathbf{e}_1+\mathbf{e}_2}\left(a_2\theta_{i+1}^{\mathbf{e}_2}\,\theta_i^{2\mathbf{e}_1} - a_1\theta_{i+1}^{2\mathbf{e}_1}\,\theta_i^{\mathbf{e}_2}\right)}{\theta_i^{\mathbf{e}_1+\mathbf{e}_2}\,\theta_{i+1}^{\mathbf{e}_1}\left(a_2\theta_i^{\mathbf{e}_2}\,\theta_{i-1}^{2\mathbf{e}_1} - a_1\theta_i^{2\mathbf{e}_1}\,\theta_{i-1}^{\mathbf{e}_2}\right)}, \tag{9.11}$$

where we omit the superscripts t for simplicity. At the second equality we have canceled all the c_i using $c_i^{\mathbf{e}_u} = c_{i+u}$. Due to (9.6), we obtain $f_i^t = 1$. Thus, using (9.4), we obtain (9.3) . ☐

9.3. **General m case.** The vertical actions e_u $(u = 1, \dots, m)$ are expressed as matrix equations:

$$Q_1^{t+\mathbf{e}_u} \cdots Q_m^{t+\mathbf{e}_u} = Q_{u+1}^t \cdots Q_m^t P(x)^k Q_1^t \cdots Q_u^t P(x)^{-k}. \tag{9.12}$$

Among the family of difference equations we will use the following ones later:

$$\prod_{j=1}^m q_{j,i}^{t+\mathbf{e}_u} = \prod_{j=u+1}^m q_{j,i}^t \cdot \prod_{j=1}^u q_{j,i-k}^t, \tag{9.13}$$

$$\sum_{j=1}^m q_{1,i}^{t+\mathbf{e}_u} \cdots q_{j-1,i}^{t+\mathbf{e}_u}\, q_{j+1,i-1}^{t+\mathbf{e}_u} \cdots q_{m,i-1}^{t+\mathbf{e}_u} = \sum_{j=1}^m q_{s(1,i)}^t \cdots q_{s(j-1,i)}^t\, q_{s(j+1,i-1)}^t \cdots q_{s(m,i-1)}^t, \tag{9.14}$$

for $u = 1, \cdots, m$. Here we define

$$s(j,i) = \begin{cases} (j+u, i) & j = 1, \dots m-u, \\ (j+u-m, i-k) & j = m-u+1, \dots, m. \end{cases}$$

Theorem 9.5. *For any $t \in \mathbb{Z}^m$, $i \in \mathbb{Z}$ and $1 \le p < r \le m$, the θ_i^t satisfy an octahedron recurrence relation,*

$$\delta_{p,r}\, \theta_{i+1}^{t+\mathbf{e}_{p-1}+\mathbf{e}_{r-1}} \theta_i^{t+\mathbf{e}_p+\mathbf{e}_r} + a_p\, \theta_{i+1}^{t+\mathbf{e}_p+\mathbf{e}_{r-1}} \theta_i^{t+\mathbf{e}_{p-1}+\mathbf{e}_r} = a_r\, \theta_{i+1}^{t+\mathbf{e}_{p-1}+\mathbf{e}_r} \theta_i^{t+\mathbf{e}_p+\mathbf{e}_{r-1}}. \tag{9.15}$$

Here $\delta_{p.r}$ is a constant given by

$$\delta_{p,r} = a_r \frac{\Theta(p_0 + \iota(A_p - A_r))\Theta(p_0 + \iota(O - P))}{\Theta(p_0 + \iota(O - A_r))\Theta(p_0 + \iota(A_p - P))},$$

and p_0 is a zero of the Riemann theta function: $\Theta(p_0) = 0$.

Proof. The proof is similar to the $m = 2$ case. We explain the outline. In the case $(P_1, P_2, P_3, P_4) = (A_r, O, P, A_p)$ of Theorem 9.1, we obtain

$$\begin{aligned} &T_1\, \Theta(z + \iota(P - A_r))\, \Theta(z + \iota(A_p - O)) + T_2\, \Theta(z + \iota(P - O))\, \Theta(z + \iota(A_p - A_r)) \\ &\qquad = T_3\, \Theta(z + \iota(P + A_p - A_r - O))\, \Theta(z) \end{aligned} \tag{9.16}$$

where T_1, T_2 and T_3 are given by (9.8), but we replace A_1 (resp. A_2) with A_p (resp. A_r). By setting $z = \beta^t + \iota((n-i-1)(O-P) + \mathcal{A}_p + \mathcal{A}_{r-1})$ at (9.16), we get

$$T_1\, \theta_{i+1}^{t+\mathbf{e}_{p-1}+\mathbf{e}_{r-1}} \theta_i^{t+\mathbf{e}_p+\mathbf{e}_r} + T_2\, \theta_{i+1}^{t+\mathbf{e}_p+\mathbf{e}_{r-1}} \theta_i^{t+\mathbf{e}_{p-1}+\mathbf{e}_r} = T_3\, \theta_{i+1}^{t+\mathbf{e}_{p-1}+\mathbf{e}_r} \theta_i^{t+\mathbf{e}_p+\mathbf{e}_{r-1}}.$$

We take $t_0 \in \mathbb{Z}^m$ and define $p_0 := -\beta^{t_0} \in \mathbb{C}^g$ in the same manner as in the proof of Lemma 9.3. Then $q_{u,n}^{t_0-\mathbf{e}_{u-1}} = 0$ holds for $u = 1, \dots, m$. From (9.13) (resp. (9.14)) of $i = n-1$ (resp. $i = n$) and $u = p-1$ or $r-1$, it follows that

$$\frac{q_{p,n-1}^{t_0-\mathbf{e}_{p-1}}}{q_{1,n-1}^{t_0}} = \frac{q_{r,n-1}^{t_0-\mathbf{e}_{r-1}}}{q_{1,n-1}^{t_0}} = 1.$$

On the other hand, by setting $z = p_0 + \iota(A_r - P)$ at (9.16), we obtain

$$\frac{T_2}{T_3} = \frac{\theta_{n-2}^{t_0+\mathbf{e}_p-\mathbf{e}_{p-1}}\,\theta_{n-1}^{t_0+\mathbf{e}_r-\mathbf{e}_{r-1}}}{\theta_{n-2}^{t_0+\mathbf{e}_r-\mathbf{e}_{r-1}}\,\theta_{n-1}^{t_0+\mathbf{e}_p-\mathbf{e}_{p-1}}} = \frac{q_{r,n-1}^{t_0-\mathbf{e}_{r-1}}}{q_{p,n-1}^{t_0-\mathbf{e}_{p-1}}} \cdot \frac{a_p}{a_r}.$$

From the above two relations, $T_2/T_3 = a_p/a_r$ follows. Finally, by setting $z = p_0+\iota(O-P)$ at (9.16) we obtain the formula of $\delta_{p,r}$. □

The following conjecture extends Proposition 9.4.

Conjecture 9.6. Suppose θ_i^t satisfy (9.15). Then $q_{j,i}^t$ defined via (9.5) satisfy all the difference equations (9.12).

We have checked the conjecture for $m \leq 3$.

10. THE TRANSPOSED NETWORK

10.1. **Transposed Lax matrix and spectral curve.**

Corresponding to $\tilde{Q}$ of (2.3), we also identify $q \in \mathcal{M}$ with an N-tuple of $n'm \times n'm$ matrices $\tilde{Q} := (\tilde{Q}_i(x))_{i\in[N]}$ as §7.1. The matrices $\tilde{Q}_i(x)$ are given in terms of the q_{ji} by

$$\tilde{Q}_{N+1-i}(x) := \tilde{P}(x) + \mathrm{diag}[q_{m,i},\ldots,q_{1,i},q_{m,i+k},\ldots,q_{1,i+k},\ldots,q_{m,i-k},\ldots,q_{1,i-k}],$$

where $\tilde{P}(x)$ is the $mn' \times mn'$ matrix:

$$\tilde{P}(x) := \begin{pmatrix} 0 & 0 & 0 & x \\ 1 & 0 & 0 & 0 \\ 0 & \ddots & \ddots & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

We define $\tilde{L}(x) := \tilde{Q}_1(x)\cdots\tilde{Q}_N(x)\tilde{P}(x)^{\bar{k}'m}$, which is another Lax matrix.

Also define a map $\tilde{\psi} : \mathcal{M} \to \mathbb{C}[x,y]$ given as a composition,

$$\tilde{Q} \mapsto \tilde{L}(x) \mapsto \det(\tilde{L}(x) - y).$$

Consequently, for $q \in \mathcal{M}$ we have two affine plane curves $\tilde{C}_{\psi(q)}$ and $\tilde{C}_{\tilde{\psi}(q)}$in $\mathbb{C}^2$, given by the zeros of $\psi(q)$ and $\tilde{\psi}(q)$ respectively. The proof of Proposition 10.1 is similar to that of Proposition 3.3.

Proposition 10.1. *The Newton polygon $N(\tilde{\psi}(q))$ is the triangle with vertices $(0, mn')$, $(m\bar{k}', 0)$ and $(m\bar{k}' + N, 0)$, where the lower hull (resp. upper hull) consists of one edge with vertices $(m\bar{k}', 0)$ and $(0, mn')$ (resp. $(m\bar{k}' + N, 0)$ to $(0, mn')$).*

Lemma 10.2. *The affine transformation*

$$\begin{pmatrix} i \\ j \end{pmatrix} \mapsto \begin{pmatrix} \bar{k}'(k+m) \\ -n'k \end{pmatrix} + \begin{pmatrix} -\bar{k}' & (1-k'\bar{k}')/n' \\ n' & k' \end{pmatrix} \begin{pmatrix} i \\ j \end{pmatrix} \tag{10.1}$$

sends integer points of $N(\psi(q))$ into integer points of $N(\tilde{\psi}(q))$.

Proof. It is easy to see that it sends the vertices correctly. By the definition of $\bar{k}'$ we know $(1-k'\bar{k}')/n'$ is an integer. Thus this transformation sends integer points to integer points. So does the inverse, as the determinant of the matrix involved is -1. □

Example 10.3. Let $(n, m, k) = (6, 3, 4)$. The two Newton polygons $N(\psi(q))$ and $N(\tilde{\psi}(q))$ are illustrated in Figure 11. Here i labels the horizontal axis and j labels the vertical axis. The dots of the same color show integer points inside the Newton polygon that get sent to each other by the transformation (10.1). The formula for the transformation in this case is



FIGURE 11.

$$\begin{pmatrix} i \\ j \end{pmatrix} \mapsto \begin{pmatrix} 14 \\ -12 \end{pmatrix} + \begin{pmatrix} -2 & -1 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} i \\ j \end{pmatrix}.$$

Proposition 10.4. *For $q \in \mathcal{M}$, the polynomials $\psi(q)$ and $\tilde{\psi}(q)$ coincide up to the monomial transformation induced by* (10.1)*. The signs of the new monomials are derived from the rule given in Theorem 3.2: the sign of $x^a y^b$ is $(-1)^{(mn'-b-1)a+b}$.*

See § 10.3 for the proof.

10.2. **Special points on the transposed curve.** We write $\tilde{f}(x, y)$ for the polynomial obtained from the fixed polynomial $f(x, y)$ in §3.3 via the affine transformation (10.1). Let $C_{\tilde{f}}$ be the smooth compactification of the affine plane curve $\tilde{C}_{\tilde{f}}$ given by $\tilde{f}(x, y) = 0$. As for C_f (Lemma 3.8), $C_{\tilde{f}}$ has a unique point $\tilde{P}$ lying over ∞. Due to this fact and Proposition 10.4, the two curves C_f and $C_{\tilde{f}}$ are isomorphic. Let $\tau : C_{\tilde{f}} \to C_f$ be the birational isomorphism, which is given by

$$(x, y) \mapsto (y^{n'} x^{k'}, x^{\frac{1-k'\bar{k}'}{n'}} y^{k'})$$

when $(x, y) \in C_{\tilde{f}} \cap (\mathbb{C}^*)^2$. Besides $\tilde{P}$, on $C_{\tilde{f}}$ we have special points $\tilde{O}_r = ((-1)^{mn'}\sigma_r, 0)$ for $r \in [N]$, and $\tilde{A}_i$ for $i \in [m]$, where near $\tilde{A}_i$ there is a local coordinate u such that

$$(x, y) \sim (u^{n'}, -(-\epsilon_i)^{\frac{1}{n'}} u^{\bar{k}'}).$$

We see that $\tau(\tilde{P}) = P$, $\tau(\tilde{A}_i) = A_i$ and $\tau(\tilde{O}_r) = O_r$.

10.3. **Proof of Proposition 10.4.** The terms contributing to a particular coefficient of the spectral curve are described by Theorem 3.2. We shall exhibit a bijection between the terms of the coefficient associated with the lattice point (i, j) inside the Newton polygon $N(\psi(q))$ and the terms contributing to the coefficient associated with the image of that point in $N(\tilde{\psi}(q))$ under the affine transformation (10.1).

An *underway path* in the network G is the mirror symmetric version of a highway path. Thus the weights of an underway path are given by Figure 5 with 0 and q_{ij} swapped. The crucial observation is that families of highway paths on the network associated with $\tilde{Q}$ are families of underway paths on the original network, parsed in the opposite direction. This is because by the construction of the transpose map between Q and $\tilde{Q}$, the corresponding toric networks are the same but are viewed from opposite sides of the torus (inside vs outside).

Now, assume we have a closed family of highway paths contributing to one of the coefficients of $\psi(q)$. Simply complement all edges that ended up on our family of closed highway paths inside the set of all edges of the network. We claim that the result can be parsed as the desired family of closed underway paths.

Indeed, the original family can be viewed as a number of horizontal "steps through" picking up a weight q_{ij} at some node, connected by intervals of staircase paths that do not pick up any weight. We can interpret our procedure as complementing used intervals of staircase paths, that is, making them not used, and vice versa. As a result, locally around each weight q_{ij} that was picked up the new path will look like what is shown in Figure 12. Thus, it will be an underway path, and it will pick up exactly such q_{ij}. In other words, the weight of the original highway family is the same as the weight of this new underway family.



FIGURE 12.

It is also easy to see that the new underway family is closed. This completes the proof.

Example 10.5. Let $(\mathfrak{a}, \mathfrak{b}, \mathfrak{c}, \mathfrak{d}) = (3, 2, 3, 2)$ as in Example 2.2. Figure 13 shows an example of a family of paths contributing to the purple term of the spectral curve as marked in Figure 11. The first step takes the complement of edges of this family inside the set of all edges of this network. The second step does not change the network or the paths, it just changes the point of view and reverses paths' directions.

11. RELATION TO THE DIMER MODEL

In this section, we give the explicit relation between the R-matrix dynamics on our toric network and cluster transformations on the honeycomb dimer on a torus. See [GK] for background on the dimer model.

11.1. **Cluster transformations on the honeycomb dimer.** The calculation in this section is the dimer analogue of the highway network computation of the geometric R-matrix (cf. [LP13, Theorem 6.2]), which was explained earlier in §4.5 (Figure 9).



FIGURE 13.

Fix a positive integer L, and consider a honeycomb bipartite graph on a cylinder as in Figure 14, where $\alpha_i, \beta_i, \gamma_i$ $(i \in \mathbb{Z}/L\mathbb{Z})$ are the weights of the faces in three cyclically consecutive rows. We write $\alpha := (\alpha_i, \beta_i, \gamma_i)_{i \in \mathbb{Z}/L\mathbb{Z}}$.



FIGURE 14. Honeycomb dimer on a cylinder

For the weights α of the honeycomb, we define a transformation R_α of α in a following way: first we split the α_1-face into two, by inserting a digon of weight -1 as the top of Figure 15. We thank R. Kenyon for explaining this operation to us.

Set the weights of the new two faces to be $-c$ and $\frac{\alpha_1}{c}$, where c is a nonzero parameter which will be determined. Let D be the quiver dual to the bipartite graph, drawn in blue in the figure. The weights of faces are to be regarded as *coefficient variables* associated to each vertex of D. With $1, \dots, L, a$ and b, we assign the vertices of D in the middle row, as depicted. Next, we apply the cluster mutations $\mu_1, \mu_2, \ldots, \mu_L$ to (D, α) in order, recursively defining ω_i $(i = 1, \ldots, L)$ by

$$\omega_1 := \frac{\alpha_1}{c}, \qquad \omega_i := \alpha_i(1 + \omega_{i-1}).$$

The condition that the digon's weight is again -1 after the L mutations gives an equation for c as $-c(1 + \omega_L) = -1$. By solving it, c is determined to be

$$c = \frac{1 - \prod_{s=1}^{L} \alpha_s}{\sum_{t=0}^{L-1} \prod_{s=1}^{t} \alpha_{1-s}},$$

and $\omega_i := \omega_i(\alpha_1, \ldots, \alpha_L)$ is obtained as

$$\omega_i(\alpha_1, \ldots, \alpha_L) = \frac{\alpha_i \sum_{t=0}^{L-1} \prod_{s=1}^{t} \alpha_{i-s}}{1 - \prod_{s=1}^{L} \alpha_s}. \tag{11.1}$$

Definition 11.1. Let R_α be the transformation of α given by

$$(\alpha_i, \beta_i, \gamma_i) \mapsto \left(\omega_{i-1}^{-1}(1 + \omega_i),\ \beta_i(1 + \omega_{i-1}^{-1})^{-1},\ \gamma_i(1 + \omega_i^{-1})^{-1}\right), \tag{11.2}$$

which is induced by the sequence of mutations $\mu_L \mu_{L-1} \cdots \mu_1$,

Due to the expression of ω_i, we see that R_α does not depend on which α_j-face we split at the first stage.

Remark 11.2. Note that our derivation of R_α is not entirely a cluster algebra computation since we began with the "digon insertion" operation, which does not have a clear cluster algebra interpretation. In an upcoming work [ILP], we plan to further clarify the cluster nature of the geometric R-matrix.

11.2. **Relation with the (n, m, k)-network.** We consider the honeycomb bipartite graph on a torus as in Figure 16. We assign each face (resp. edge) with a weight x_{ji} (resp. q_{ji} or 1), satisfying the periodicity conditions $x_{j,i+n} = x_{j,i}$ and $x_{m+j,i} = x_{j,i-k}$ (resp. $q_{j,i+n} = q_{j,i}$ and $q_{m+j,i} = q_{j,i-k}$). In the figure we omit weights that are equal to 1. The x_{ji} and q_{ji} are related by

$$x_{j,i} = \frac{q_{j,i+1}}{q_{j+1,i}} \ (j = 1, \ldots, m-1), \qquad x_{m,i} = \frac{q_{m,i+1}}{q_{1,i-k}}, \tag{11.3}$$

hence the product of all the x_{ji} is equal to 1.

Let $\mathcal{X} \simeq \mathbb{C}^{mn}$ be a space of the weights of faces whose coordinates are given by $\underline{x} := (x_{ji})_{j \in \mathbb{Z}/m\mathbb{Z},\ i \in \mathbb{Z}/n\mathbb{Z}}$. Let ρ be an embedding map from $\mathbb{C}(\mathcal{X})$ to $\mathbb{C}(\mathcal{M})$ given by (11.3).

We define three types of actions $\bar{R}_j$, $\tilde{R}_i$ and $\hat{R}_i$ on $\mathcal{X}$, corresponding to the three directions that the honeycomb bipartite graph can be arranged into rows. For $j = 1, \ldots, m$, define $\bar{x}(j) := (x_{j,i}, x_{j-1,i}, x_{j+1,i})_{i \in \mathbb{Z}/n\mathbb{Z}}$. Let $\bar{R}_j$ be the action on $\mathcal{X}$ induced by $R_{\bar{x}(j)}$ (11.2) with $L = n$. More precisely, $R_j(\underline{x}) = \underline{x}'$ is given by

$$x'_{li} = \begin{cases} \omega_{i-1}^{-1}(1 + \omega_i) & l = j, \\ x_{j-1,i}(1 + \omega_{i-1}^{-1})^{-1} & l = j-1, \\ x_{j+1,i}(1 + \omega_i^{-1})^{-1} & l = j+1, \\ x_{li} & \text{otherwise}, \end{cases}$$


γ_{L-1}
γ_L
γ_1
γ_2
L
a
b
1
2
α_L
$-c$
-1
$\frac{\alpha_1}{c}$
α_2
β_L
β_1
β_2
β_3
$\downarrow\ \mu_1$
γ_{L-1}
γ_L
$\gamma_1(1+\frac{c}{\alpha_1})^{-1}$
γ_2
L
a
b
1
2
α_L
$-c$
$-(1+\frac{\alpha_1}{c})$
$\frac{c}{x_1}$
$\alpha_2(1+\frac{\alpha_1}{c})$
β_L
β_1
$\beta_2(1+\frac{c}{\alpha_1})^{-1}$
β_3
$\downarrow\ \mu_L\cdots\mu_2$
$\gamma_{L-1}(1+\frac{1}{\omega_{L-1}})^{-1}$
$\gamma_L(1+\frac{1}{\omega_L})^{-1}$
$\gamma_1(1+\frac{1}{\omega_1})^{-1}$
$\gamma_2(1+\frac{1}{\omega_2})^{-1}$
$-c(1+\omega_L)$
$L-1$
L
a
b
1
$-(1+\omega_1)$
$\frac{1+\omega_L}{\omega_{L-1}}$
$\frac{1}{\omega_L}$
$\frac{1+\omega_2}{\omega_1}$
$\beta_L(1+\frac{1}{\omega_{L-1}})^{-1}$
$\beta_1(1+\frac{1}{\omega_L})^{-1}$
$\beta_2(1+\frac{1}{\omega_1})^{-1}$
$\beta_3(1+\frac{1}{\omega_2})^{-1}$

Figure 15. Honeycomb dimer model on a cylinder.

with $\omega_i := \omega_i(x_{j,1},\ldots,x_{j,n})$ (11.1). In a similar way, for $i = 1,\ldots,N$, define $\tilde{x}(i) := (x_{m-j,N-i}, x_{m-j,N-i-1}, x_{m-j,N-i+1})_{j\in\mathbb{Z}/mn'\mathbb{Z}}$ and let $\tilde{R}_i$ be the action on $\mathcal{X}$ given by $R_{\tilde{x}(i)}$ (11.2) with $L = mn'$. For $i = 1,\ldots,M$, define $\hat{x}(i) := (x_{j,i+1-j}, x_{j,i+2-j}, x_{j,i-j})_{j\in\mathbb{Z}/\frac{mn}{M}\mathbb{Z}}$, and let $\hat{R}_i$ be the action on $\mathcal{X}$ given by $R_{\hat{x}(i)}$ (11.2) with $L = \frac{mn}{M}$.



FIGURE 16. (n, m, k)-dimer

Recall that we also have three types of actions on $\mathcal{M}$ given by the elements s_j ($j = 0, 1, \dots, m-1$) of the extended symmetric group W, the elements $\tilde{s}_i$ ($i = 0, 1, \dots, N-1$) of the extended symmetric group $\tilde{W}$, and the snake path action T_s ($s = 1, \dots, M$).

Proposition 11.3.

(i) *The actions s_j and $\tilde{s}_i$ are compatible with the actions $\bar{R}_j$ and $\tilde{R}_j$ respectively: for $\underline{x} \in \mathcal{X}$, we have*

$$\rho \circ \bar{R}_j(\underline{x}) = s_j^* \circ \rho(\underline{x}), \quad j \in \mathbb{Z}/m\mathbb{Z}, \tag{11.4}$$

$$\rho \circ \tilde{R}_i(\underline{x}) = \tilde{s}_i^* \circ \rho(\underline{x}), \quad i \in \mathbb{Z}/N\mathbb{Z}. \tag{11.5}$$

(ii) *For $\underline{x} \in \mathcal{X}$, the action $\hat{R}_i$ satisfies*

$$\rho \circ \hat{R}_i(\underline{x}) = \rho(\underline{x}), \quad i = 1, \dots, M. \tag{11.6}$$

(iii) *For $\underline{x} \in \mathcal{X}$, the snake path action T_i^* satisfies*

$$T_i^* \circ \rho(\underline{x}) = \rho(\underline{x}), \quad i = 1, \dots, M.$$

Proof. (i) To show (11.4), it is enough to prove the $j = 1$ case. We write E_i for the energy $E(P^{-i}Q_1P^i, P^{-i}Q_2P^i)$. The operator s_1 acts on $\mathcal{M}$ as $s_1(q) = q'$:

$$q'_{1,i} = q_{2,i}\,\frac{E_i}{E_{i-1}}, \qquad q'_{2,i} = q_{1,i}\,\frac{E_{i-1}}{E_{j,i}}, \tag{11.7}$$

and the other q_{ji} do not change. By definition, $R_1(\underline{x}) = \underline{x}'$ is obtained as

$$x'_{ji} = \begin{cases} \omega_{i-1}^{-1}(1+\omega_i) & j = 1, \\ x_{m,i+k}(1+\omega_{i-1}^{-1})^{-1} & j = m, \\ x_{2,i}(1+\omega_i^{-1})^{-1} & j = 2, \\ x_{ji} & \text{otherwise.} \end{cases}$$

where $\omega_i := \omega_i(x_{1,1}, \dots, x_{1,n})$. On the other hand, by direct computation we obtain

$$\rho(\omega_i) = \frac{q_{1,i+1} E_i}{\prod_{s=1}^n q_{2,s} - \prod_{s=1}^n q_{1,s}}, \qquad \rho(1+\omega_i) = \frac{q_{2,i+1} E_{i+1}}{\prod_{s=1}^n q_{2,s} - \prod_{s=1}^n q_{1,s}}.$$

Thus we get

$$\begin{aligned} \rho\left(\omega_{i-1}^{-1}(1+\omega_i)\right) &= \frac{q_{2,i+1}E_{i+1}}{q_{1,i}E_{i-1}} = s_1^* \left(\frac{q_{1,i+1}}{q_{2,i}}\right) = s_1^* \circ \rho(x_{1,i}), \\ \rho\left(x_{m,i+k}(1+\omega_{i-1}^{-1})^{-1}\right) &= \frac{q_{m,i+k+1}E_{i-1}}{q_{2,i}E_i} = s_1^* \left(\frac{q_{m,i+k+1}}{q_{1,i}}\right) = s_1^* \circ \rho(x_{m,i+k}), \\ \rho\left(x_{2,i}(1+\omega_i^{-1})^{-1}\right) &= \frac{q_{1,i+1}E_i}{q_{3,i}E_{i+1}} = s_1^* \left(\frac{q_{2,i+1}}{q_{3,i}}\right) = s_1^* \circ \rho(x_{2,i}), \end{aligned}$$

and (11.4) follows.

We can prove (11.5) in a similar manner, by replacing Q_i with $\tilde{Q}_i$, P with $\tilde{P}$, and so on.

(ii) Again, it is enough to prove the case of $i = 1$. For simplicity, we write L for $\frac{mn}{M}$, and set $x_j := x_{j,2-j}$, $q_j := q_{j,3-j}$ for $j \in \mathbb{Z}/L\mathbb{Z}$. From (11.3), it follows that

$$\rho(x_j) = \frac{q_j}{q_{j+1}}, \qquad \rho\left(\prod_{j=1}^L x_j\right) = 1. \tag{11.8}$$

Define $\omega_j = \omega_j(x_1, \dots, x_L)$ for $j = 1, \dots, L$. Using the definition of ω_j and (11.8), we obtain

$$\rho\left(\frac{1+\omega_j}{\omega_{j-1}}\right) = \frac{q_j}{q_{j+1}}, \qquad \rho\left(\frac{\omega_j}{1+\omega_j}\right) = 1, \tag{11.9}$$

by the following calculation:

$$\begin{aligned} \frac{1+\omega_j}{\omega_{j-1}} &= \frac{1 - \prod_{s=1}^L x_s + x_i \sum_{t=0}^{L-1} \prod_{s=1}^t x_{i-s}}{x_{i-1} \sum_{t=0}^{L-1} \prod_{s=1}^t x_{i-1-s}} \\ &\xmapsto{\rho} \frac{1 - 1 + \sum_{t=0}^{n-1} \frac{q_{i-t}}{q_{i+1}}}{\sum_{t=1}^n \frac{q_{i-t}}{q_i}} = \frac{q_i}{q_{i+1}}, \\ \frac{\omega_j}{1+\omega_j} &= \frac{x_i \sum_{t=0}^{L-1} \prod_{s=1}^t x_{i-s}}{1 - \prod_{s=1}^L x_s + x_i \sum_{t=0}^{L-1} \prod_{s=1}^t x_{i-s}} \xmapsto{\rho} 1. \end{aligned}$$

Thus we have $\rho \circ R_{\hat{x}(1)}(\hat{x}(1)) = \rho(\hat{x}(1))$, and (11.6) follows.

(iii) The snake path action T_s changes $q_{j,i}$ and $q_{j',i'}$ in the same way if $i + j \equiv i' + j'$ mod M. This condition is satisfied by $q_{j,i+1}$ and $q_{j+1,i}$, then the change is cancelled in $\rho(x_{j,i})$. Thus we see that $T_s \circ \rho(x_{j,i}) = \rho(x_{j,i})$. ☐

REFERENCES

[BK] A. Berenstein and D. Kazhdan, Geometric and unipotent crystals, Geom. Funct. Anal., Special Volume, Part I, 188–236 (2000).

[Eti] P. Etingof, Geometric crystals and set-theoretical solutions to the quantum Yang-Baxter equation, Comm. Algebra, 31, no. 4, 1961–1973 (2003).

[Fay] J. D. Fay, *Theta functions on Riemann surfaces*, Lecture notes in Math., **352** (Springer-Verlag, Berlin-New York, 1973).

[FM] V. V. Fock and A. Marshakov, Loop groups, Clusters, Dimers and Integrable systems, arXiv:1401.1606.

[GSTV] M. Gekhtman, M. Shapiro, S. Tabachnikov, A. Vainshtein, Integrable cluster dynamics of directed networks and pentagram maps, arXiv:1406.1883.

[GK] A. B. Goncharov and R. Kenyon, Dimers and cluster integrable systems, Ann. Sci. Éc. Norm. Supér. (4) 46, no. 5, 747–813 (2013).

[HTI] R. Hirota, S. Tsujimoto and T. Imai, Difference scheme of soliton equations, in Future directions of nonlinear dynamics in physical and biological systems (Lyngby, 1992), NATO Adv. Sci. Inst. Ser. B Phys., 312, 7–15 (Plenum Press, New York, 1993).

[IKT] R. Inoue, A. Kuniba and T. Takagi, Integrable structure of box-ball systems: crystal, Bethe ansatz, ultradiscretization and tropical geometry, J. Phys. A **45**, no. 7, 073001, 64 pp (2012).

[IT] R. Inoue and T. Takenawa, Tropical spectral curves and integrable cellular automata, Int. Math. Res. Not. IMRN **27** Art ID. rnn019, 27 pp (2008);

———, A tropical analogue of Fay's trisecant identity and an ultradiscrete periodic Toda lattice, Comm. Math. Phys., 289, 995–1021 (2009).

[ILP] R. Inoue, T. Lam, and P. Pylyavskyy, in preparation.

[I08] S. Iwao, Solution of the generalized periodic discrete Toda equation, J. Phys. A 41, no. 11, 115201, 15 pp (2008).

[I10] S. Iwao, Solution of the generalized periodic discrete Toda equation. II. Theta function solution, J. Phys. A 43, no. 15, 155208, 9 pp (2010).

[I09] S. Iwao, Linearisation of the (M, K)-reduced non-autonomous discrete periodic KP equation, arXiv:0912.3333.

[Kir] A.N. Kirillov, Introduction to tropical combinatorics, in "Physics and Combinatorics 2000" (Eds. A. N. Kirillov and N. Liskova), Proceedings of the Nagoya 2000 International Workshop, 82–150, World Scientific, 2001.

[Kho] A. Khovanskii, Three Lectures on Newton Polyhedra (Newton polyhedrons and Singularities). RIMS Kokyuroku, 1233, 1–17 (2001).

[KKMMNN] S.-J. Kang, M. Kashiwara, K.C. Misra, T. Miwa, T. Nakashima, and A. Nakayashiki, Affine crystals and vertex models, in Infinite analysis Part A (Kyoto 1991), 449–484, *Adv. Ser. Math. Phys.*, **16**, World Sci. Publishing, River Edge, NJ, 1992.

[KN1] M. Kashiwara, T. Nakashima, and M. Okado, Affine geometric crystals and limit of perfect crystals. (English summary) *Trans. Amer. Math. Soc.* 360, no. 7, 3645–3686 (2008).

[KN2] M. Kashiwara, T. Nakashima, and M. Okado, Tropical R maps and affine geometric crystals, *Represent. Theory* 14, 446–509 (2010).

[KNY] K. Kajiwara, M. Noumi, and Y. Yamada, Discrete Dynamical Systems with $W(A^{(1)}_{m-1} \times A^{(1)}_{n-1})$ Symmetry, Lett. Math. Phys. 60, no. 3, 211–219 (2002).

[KT] T. Kimijima and T. Tokihiro, Initial-value problem of the discrete periodic Toda equation and its ultradiscretization, Inverse Problems, textbf18, no. 6, 1705–1732 (2002).

[Las] A. Lascoux, Double crystal graphs. Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), 95–114, Progr. Math., 210, Birkhäuser Boston, Boston, MA, 2003.

[LP11] T. Lam and P. Pylyavskyy, Affine geometric crystals in unipotent loop groups, Representation Theory 15, 719–728 (2011).

[LP12a] T. Lam and P. Pylyavskyy, Total positivity in loop groups, I: Whirls and curls. Adv. Math. 230, no. 3, 1222–1271 (2012).

[LP12b] T. Lam and P. Pylyavskyy, Inverse problem in electrical cylindrical networks, SIAM J. of Applied Math., 72, 767–788 (2012).

[LP13] T. Lam and P. Pylyavskyy, Crystals and total positivity on orientable surfaces, Selecta Math. (N.S.) **19**, no. 1, 173–235 (2013).

[MIT] J. Mada, M. Idzumi and T. Tokihiro, Path description of conserved quantities of generalized periodic box-ball systems, J. Math. Phys. **46**, no. 2, 022701, 19 pp (2005).

[Rho] B. Rhoades, Cyclic sieving, promotion, and representation theory, J. Combin. Theory Ser. A 117, no. 10, no. 1, 38–76 (2010).

[Sch] M.-P. Schützenberger, Promotion des morphismes d'ensembles ordonnes, Discrete Math. no.2, 73–94 (1972).

[Spe] D. Speyer, Perfect matchings and the octahedron recurrence, J. Algebraic Combin. 25, no. 3, 309–348 (2007).

[TS] D. Takahashi and J. Satsuma, A soliton cellular automaton, J. Phys. Soc. Japan **59**, no. 10, 3514–3519 (1990).

[vMM] P. van Moerbeke and D. Mumford, The spectrum of difference operators and algebraic curves, Acta Math. **143** (1-2) 94–154 (1979).

Rei Inoue, Department of Mathematics and Informatics, Faculty of Science, Chiba University, Chiba 263-8522, Japan.
E-mail address: `reiiy@math.s.chiba-u.ac.jp`

Thomas Lam, Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA.
E-mail address: `tfylam@umich.edu`

Pavlo Pylyavskyy, School of Mathematics, University of Minnesota, Minneapolis, MN 55414, USA.
E-mail address: `ppylyavs@umn.edu`